arXiv:cond-mat/9509090v1 15 Sep 1995

NIM-TH 95-11

Talk at the second pacific winter School
in Theoretical Physics. Korea, 1995

Introduction to *Yangian* in Physics

Mo-Lin Ge†, Kang Xue‡ and Yiwen Wang†

†*Theoretical Physics Division, Nankai Institute of Mathematics,*
Tianjin 300071, People's Republic of China
‡*Physics Department, Northeast Normal University,*
Changchun 130024, People's Republic of China

Abstract

This is an introduction to the physical pictures of *Yangian* symmetry. All the discussions are based on the RTT relations which have been known to be related to the Hamiltonian formulations for quantum integrable systems. The explicit calculations of *Yangian* associated with $gl(2)$ are given through solving the RTT relations. It contains the Drinfeld's $Y(sl(2))$, $Y(sl(n))$ and Heisenberg type of Hopf algebras. Examples are emphasized to show the physical pictures, such as Hubbard model, Haldane-Shastry model, Long-range interaction models and Goryachev-Chaplygin(G-C) gyrostat. The example of the trigonometric extension of G-C gyrostat are also shown that leads to the non-commutative geometry naturally on the basis of the truncated q-affine algebra.

§1 Introduction

An integrable system is said to be solved if the equation of motion is solved or all the conserved quantities are found. From the point of view of Nöther theorem equations of motion and conserved quantities co-exist in connecting with certain symmetries of the $\mathcal{L}$agrangian. In quantum mechanics, to obtain the spectrum we can either solve the Schödinger equation or find all the symmetries in the system. The typical example is the Hydrogen atom which possesses another SU(2) invariance due to the Runge-Lenz vector besides the well-known SU(2) associated with the spherically symmetric potential U(r). With the given SO(4)=SU(2)⊗SU(2)) symmetry the spectrum of a Hydrogen atom can easily be obtained without the explicit form of the wave function satisfying Schrödinger equation [1]. In this example the family of the symmetries consist in the two SU(2) that commute with each other, namely, the two conserved quantum operators corresponding to the symmetries form a commuting family. This simple example provides a representation of completely quantum integrable systems.

To extend the above analysis to nonlinear models it should be noticed that a quantum many-body problem in the first quantization form with conserved particle number N is, in principle, equivalent to the corresponding quantum field theory(second quantization form). The typical evidence is the equivalence between δ-interaction model with the Hamiltonian

$$H = -\sum_{i=1}^{N} \partial_i^2 + 2c \sum_{i \neq j} \delta(x_i - x_j) \tag{1.1}$$

and the nonlinear Schrödinger equation [2]:

$$\hat{H} = \int dx \left(\frac{1}{2} \partial_x \phi \partial_x \phi^* + 2c(\phi\phi^*)^2 \right) \tag{1.2}$$

where $\phi = \phi(x,t)$ stands for the second quantized operator.

In solving a many-body problem we meet also two approaches. The first one is to solve the Schrödinger equation directly. It is the well-known Bethe-Anastz methods [3]. The second approach is to find the commuting family of conserved quantities consisting of quantum operators, that is known as the Quantum Inverse Scattering Methods(QISM) [4]. Although the Bethe-Ansatz approach is more powerful in solving the 1-dim. many-body problems, the QISM provides a model-independently systematic method to deal with the (1+1) dimensional integrable QFT [4]. In this approach the RTT relation plays the central role. For a wide class of models it takes the form [5]:

$$R(u-v)(T(u) \otimes I)(I \otimes T(v)) = (I \otimes T(v))(T(u) \otimes I)R(u-v) \tag{1.3}$$

where $T(u)$ stands for the quantum transfer matrix and u and v the spectral parameters. In general,

$$T(u) = ||T_{ab}(u)||_{a,b=1}^{N} \tag{1.4}$$

and R is then N^2 by N^2 c-numbered matrix. In eq.(1.4) each element $T_{ab}(u)$ is u-dependent quantum operator. For this reason the space spanned by the indices $a, b, \ldots$ is called "auxiliary space", whereas the Hilbert space spanned by T_{ab} itself is called "quantum space".

Defining

$$\overset{(1)}{T}(u) = T(u) \otimes I, \quad \overset{(2)}{T}(v) = I \otimes T(v) \tag{1.5}$$

the eq.(1.5) can be recast into the form [5]

$$R(u-v) \overset{(1)}{T}(u) \overset{(2)}{T}(v) = \overset{(2)}{T}(v) \overset{(1)}{T}(u) R(u-v) \tag{1.6}$$

where each element of T is a quantum operator so that the ordering for $\overset{(1)}{T}$ and $\overset{(2)}{T}$ is very important. Noting that

$$(\overset{(1)}{T}(u) \overset{(2)}{T}(v))_{ij,kl} = (T(u) \otimes I)_{ij,mn} (I \otimes T(v)_{mn,kl} = (T(u) \otimes T(v))_{ij,kl}$$

the component form of eq. (1.6) is

$$R(u-v)_{ij,ms} (T(u) \otimes T(v))_{ms,kl} = (T(v) \otimes T(u))_{ji,sr} R(u-v)_{rs,kl}$$

that can be rewritten in the form

$$\check{R}(u-v)_{ij,rs} (T(u) \otimes T(v))_{rs,kl} = (T(v) \otimes T(u))_{ij,rs} \check{R}(u-v)_{rs,kl}$$

i.e.,

$$\check{R}(u-v)(T(u) \otimes T(v)) = (T(v) \otimes T(u)) \check{R}(u-v) \tag{1.7}$$

where

$$\check{R}(u) = PR(u) \quad \text{or} \quad \check{R}_{ij,kl} = R_{ji,kl} \tag{1.8}$$

From eq. (1.6) or eq. (1.7) it follows:

(1) Because $\check{R}(u)$ is non-singular matrix:

$$T(u) \otimes T(v) = \check{R}^{-1}(u-v)(T(v) \otimes T(u)) \check{R}(u-v) \tag{1.9}$$

hence, by taking the trace of both of sides we have

$$[\mathrm{tr}T(u), \ \mathrm{tr}T(v)] = 0 \tag{1.10}$$

where the commutation relation is taken over the quantum space. By taking the expansion in terms of u and v

$$\mathrm{tr}T(u) = \sum u^{-n} J^{(n)} \tag{1.11}$$

we get

$$[J^{(n)}, J^{(m)}] = 0 \tag{1.12}$$

i.e., there exists an infinite set of conserved quantum operators $J^{(n)}$.

(2) In eq. (1.6) the R-matrix gives rise to the exchange between the space 1 and space 2, we can denote the R by R_{12}. Similarly it can be introduce R_{ij} through

$$R_{ij} \overset{(i)}{T}\overset{(j)}{T} = \overset{(j)}{T}\overset{(i)}{T} R_{ij}$$

then

$$R_{12}R_{13}R_{23} \overset{1}{T}\overset{2}{T}\overset{3}{T} = \overset{3}{T}\overset{2}{T}\overset{1}{T} R_{12}R_{13}R_{23}$$

and

$$R_{23}R_{13}R_{12} \overset{1}{T}\overset{2}{T}\overset{3}{T} = \overset{3}{T}\overset{2}{T}\overset{1}{T} R_{23}R_{13}R_{12}$$

that lead to

$$[R_{12}R_{13}R_{23}(R_{23}R_{13}R_{23})^{-1}, \overset{3}{T}\overset{2}{T}\overset{1}{T}] = 0$$

where the first term in the commutator should be unity because its determinant is unity. Thus base on the associativity we have the Yang-Baxter equation(YBE):

$$R_{12}(u_{12})R_{13}(u_{13})R_{23}(u_{23}) = R_{23}(u_{23})R_{13}(u_{13})R_{12}(u_{12}) \ (u_{ij} = u_i - u_j) \tag{1.13}$$

YBE is a set of algebraic equations. The component form of YBE reads:

$$R^{ij}_{rs}(u_{12})R^{rk}_{lp}(u_{13})R^{sp}_{lm}(u_{23}) = R^{jk}_{sp}(u_{23})R^{ip}_{rn}(u_{13})R^{rs}_{lm}(u_{12}) \tag{1.14}$$

where the repeated indices mean summation.

Now the standard procedure to deal with a Y-B system is:

1. Solving the YBE to find a R-matrix satisfying eq. (1.14).
2. Finding the commutation relations for T_{ab} by solving the RTT relation eq. (1.6) or eq. (1.7).
3. Making a physical realization of T_{ab}.
4. Substituting the T_{ab} into tr$T$ or other conserved quantities that commute with tr$T$ to find the Hamiltonian.
5. Following the QISM to find “spectrum”.

The step 2 is crucial. The RTT relation can give rise to new types of symmetries associated with nonlinear interaction other than the Lie algebraic structures for the linear models. The simple examples of R-matrices satisfying YBE is given below:

(a) The simplest rational form [6]:

$$R(u) = u + P, \quad \check{R}(u) = uP + I \tag{1.15}$$

where P is permutation operator $P(a \otimes b) = (b \otimes a)$. For a rational R-matrix we have

$$T(u) = \sum_{n=0}^{\infty} u^{-n} T^{(n)} \tag{1.16}$$

i.e., the expansion of the transfer matrix is taken in the "half" axis of the spectral parameter u.

(b) The simplest trigonometric form [7]:

$$\begin{aligned}\check{R}(u) &= \begin{bmatrix} xq - x^{-1}q^{-1} & & & \\ & q - q^{-1} & x - x^{-1} & \\ & x - x^{-1} & q - q^{-1} & \\ & & & xq - x^{-1}q^{-1} \end{bmatrix} \\ &= \begin{bmatrix} \sinh(u+\gamma) & & & \\ & \sinh\gamma & \sinh u & \\ & \sinh u & \sinh\gamma & \\ & & & \sinh(u+\gamma) \end{bmatrix}\end{aligned} \tag{1.17}$$

where $x = e^u, \ q = e^\gamma$, or

$$R(u) = \begin{bmatrix} \sinh(u+\gamma) & & & \\ & \sinh u & \sinh\gamma & \\ & \sinh\gamma & \sinh u & \\ & & & \sinh(u+\gamma) \end{bmatrix} \tag{1.18}$$

that corresponds to the 6-vertex model. Obviously by $u \to \gamma u$ and let $\gamma \to 0$, eq. (1.17) tends to

$$\check{R}(u) \to \gamma(I + uP).$$

Noting that the overall constant factor to R-matrix can always be dropped. Therefore, eq. (1.15) is nothing but the rational limit of eq. (1.18).

(c) The simplest elliptic solution reads [8].

$$R(u) = \begin{bmatrix} a(u) & & & d(u) \\ & b(u) & c(u) & \\ & c(u) & b(u) & \\ d(u) & & & a(u) \end{bmatrix} \tag{1.19}$$

where

$$\begin{aligned}
a(u) &= \theta_0(\eta)\theta_0(u)\theta_1(u+\eta) \\
b(u) &= \theta_0(\eta)\theta_1(u)\theta_0(u+\eta) \\
c(u) &= \theta_1(\eta)\theta_0(u)\theta_0(u+\eta) \qquad (1.20) \\
d(u) &= \theta_1(\eta)\theta_1(u)\theta_1(u+\eta) \\
\theta_0(u) &= \prod_{n=1}^{\infty}(1-2p^{(n-1)/2}\cos 2\pi u + p^{2n-1})(1-p^n) \\
\theta_1(u) &= 2p^{\frac{1}{8}}\sin \pi u \prod_{n=1}^{\infty}(1-2p^n\cos 2\pi u + p^{2n})(1-p^n) \qquad (1.21)
\end{aligned}$$

with η and p being free parameters, i.e., R-matrix given by eq. (1.18) is double periodic. When $p \approx 0$, $\theta_0(u) \approx 2\sin^2(\pi u)$, $\theta_1(u) \approx 2p^{\frac{1}{8}}\sin \pi u$, we have $a(u) \approx f(p)\sin \pi(u+\eta)$, $b(u) \approx f(p)\sin \pi u$, $c(u) \approx f(p)\sin \pi\eta$ and $d(u) \approx 0$. Hence, eq.(1.19) $\longrightarrow$ eq.(1.18), namely, it is reduced to the trigonometric form.

The types (a), (b) and (c) correspond to the vertex models. Beyond them there is the fourth type of R-matrix [9]:

(d) Chiral Potts model

This type of solution is expressed in terms of functions $W_{pq}(a-b)$ and $\overline{W}_{pq}(a-b)$ that satisfy

$$\begin{aligned}
&\sum_d W_{pr}(a-d)\overline{W}_{qr}(b-d)\overline{W}_{pq}(d-c) \\
=\ & R_{pqr}\overline{W}_{pr}(b-c)W_{qr}(a-c)W_{pq}(a-b) \qquad (1.22)
\end{aligned}$$

where W and $\overline{W}$ can be illustrated in terms of the diagrams



$W_{pq}(a-b)$



$\overline{W}_{pq}(a-b)$

Eq.(1.22) can be solved:

$$\begin{aligned}
g_{pq}(n) &= \frac{W_{pq}(n)}{W_{pq}(0)} = \prod_{j=1}^{n}\frac{d_p b_q - a_p c_q \omega^j}{b_p d_q - c_p a_q \omega^j} \qquad (1.23) \\
\overline{g}_{pq}(n) &= \frac{\overline{W}_{pq}(n)}{\overline{W}_{pq}(0)} = \prod_{j=1}^{n}\frac{\omega a_p d_q - d_p a_q \omega^j}{c_p b_q - b_p c_q \omega^j} \qquad (1.24) \\
\omega &= e^{\frac{2\pi i}{N}}
\end{aligned}$$

and

$$R_{pqr} = \frac{f_{pq} f_{qr}}{f_{qr}} \tag{1.25}$$

$$f_{pq} = \left[\frac{\prod_{m=1}^{N} (\sum_{k=1}^{N} \omega^{mk} \overline{W}_{pq}(k))}{\prod_{m=1}^{N} W_{pq}(m)} \right]^{\frac{1}{N}} \tag{1.26}$$

The self-duality conditions:

$$c_p = d_p = 1, \quad a_p^N + b_p^N = 0 \tag{1.27}$$

simplify the solution to

$$g_{pq}(n) = \prod_{j=1}^{n} \frac{b_q - a_p \omega^j}{b_p - a_q \omega^j} \tag{1.28}$$

$$\overline{g}_{pq}(n) = \prod_{j=1}^{n} \frac{\omega a_p - a_q \omega^j}{b_q - b_p \omega^j} \tag{1.29}$$

where a_p, b_p and ω^j are all parameters. Especially, a_p and b_p play the role as the spectral parameter u in the vertex models.

The main goal of this lecture is to give a physical understanding on the consequences of RTT relations for R-matrices are rational and trigonometric. We shall first calculate the explicit commutation relations satisfied by T_{ab} $(a, b = 1, 2)$ with the R-matrix given by the rational form eq. (1.15), then introduce the *Yangian* symmetry given by Drinfeld [10]. We try to give a phyiscal interpretation of Drinfeld's theory. Next we would like to discuss variety of physical realizations of *Yangian* and point out how the *Yangian* symmetry is used to construct Hamiltonian for a quantum integrable system. As an example, the long-range interaction model for sl(n) will be discussed. The trigonometric solution of R-matrix eq. (1.18) gives rise to the quantum algebra. we shall explain how to find a q-deformed model based on the truncated affine quantum algebra. For reader's convenience we would like make the paper self-contained.

§2 RTT Relation and Y(sl(2))

§2.1 RTT and its Commutation Relations

For the given sl(2) R-matrix shown by eq. (1.15) the permutation P takes the form

$$P = \begin{bmatrix} 1 & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & 1 \end{bmatrix} \tag{2.1}$$

Setting the transfer matrix

$$T(u) = \left[\begin{array}{cc} T_{11}(u) & T_{12}(u) \\ T_{21}(u) & T_{22}(u) \end{array} \right] = \sum_{n=0}^{\infty} u^{-n} T^{(n)} \tag{2.2}$$

where

$$T_{ab}(u) = \sum_{n=0}^{\infty} u^{-n} T_{ab}^{(n)} \quad (a, b = 1, 2) \tag{2.3}$$

and substituting eqs. (1.15) and (2.2) into eq. (1.7) we find the relations satisfied by $T_{ab}(u)$ $(a, b, c, d = 1, 2)$:

$$(u - v)[T_{bc}(u), T_{ad}(v)] + T_{ac}(u)T_{bd}(v) - T_{ac}(v)T_{bd}(u) = 0 \tag{2.4}$$

that can be expanded in terms of u and v through eq.(2.3) $(m, n = 0, 1, \ldots, \infty)$

$$[T_{ab}^{(0)}, T^{(n)}] = 0 \tag{2.5}$$

$$[T_{bc}^{(n+1)}, T_{ad}^{(m)}] - [T_{bc}^{(n)}, T_{ad}^{(m+1)}] + T_{ac}^{(n)} T_{bd}^{(m)} - T_{ac}^{(m)} T_{bd}^{(n)} = 0 \tag{2.6}$$

eqs. (2.5) and (2.6) have the following independent relations $(a, b, c, d = 1, 2)$

$$[T_{ab}^{(m)}, T_{ab}^{(n)}] = 0, \quad [T_{ab}^{(m)}, T_{cd}^{(n)}] = [T_{ab}^{(n)}, T_{cd}^{(m)}] \tag{2.7}$$

$$\left\{ \begin{array}{l} [T_{11}^{(n+1)}, T_{12}^{(m)}] - [T_{11}^{(n)}, T_{12}^{(m+1)}] + T_{11}^{(n)} T_{12}^{(m)} - T_{11}^{(m)} T_{12}^{(n)} = 0 \\ [T_{11}^{(n+1)}, T_{21}^{(m)}] - [T_{11}^{(n)}, T_{21}^{(m+1)}] + T_{21}^{(n)} T_{11}^{(m)} - T_{21}^{(m)} T_{11}^{(n)} = 0 \\ [T_{22}^{(n+1)}, T_{12}^{(m)}] - [T_{22}^{(n)}, T_{12}^{(m+1)}] + T_{12}^{(n)} T_{22}^{(m)} - T_{12}^{(m)} T_{22}^{(n)} = 0 \\ [T_{22}^{(n+1)}, T_{21}^{(m)}] - [T_{22}^{(n)}, T_{21}^{(m+1)}] + T_{22}^{(n)} T_{21}^{(m)} - T_{22}^{(m)} T_{21}^{(n)} = 0 \end{array} \right. \tag{2.8}$$

$$\left\{ \begin{array}{l} [T_{11}^{(n+1)}, T_{22}^{(m)}] - [T_{11}^{(n)}, T_{22}^{(m+1)}] + T_{21}^{(n)} T_{12}^{(m)} - T_{21}^{(m)} T_{12}^{(n)} = 0 \\ [T_{12}^{(n+1)}, T_{21}^{(m)}] - [T_{12}^{(n)}, T_{21}^{(m+1)}] + T_{22}^{(n)} T_{11}^{(m)} - T_{22}^{(m)} T_{11}^{(n)} = 0 \end{array} \right. \tag{2.9}$$

Obviously, $T_{ab}^{(0)}$ must be a c-numbered matrix. Because the form of RTT relation is invariant under a similar transformation, without loss of generality two possible forms of matrix $T^{(0)}$

$$\text{(A)} \quad T^{(0)} = \left[\begin{array}{cc} 1 & 0 \\ 0 & \mu \end{array} \right] \quad (\mu \neq 0) \tag{2.10}$$

and

$$\text{(B)} \quad T^{(0)} = \left[\begin{array}{cc} 1 & 0 \\ 0 & 0 \end{array} \right] \tag{2.11}$$

are allowed.

§2.2 $Y(sl(2))$ Symmetry

In this section only (A) is taken into account. In this case eqs. (2.7)-(2.9) are reduced to the following independent set of relations:

$$[T_{ab}^{(m)}, T_{ab}^{(n)}] = 0, \quad [T_{ab}^{(m)}, T_{cd}^{(n)}] = [T_{ab}^{(n)}, T_{cd}^{(m)}] \ (n, m \geq 1) \tag{2.12}$$

$$\begin{cases} [T_3^{(n)}(\mu), T_{12}^{(1)}] = 2\mu T_{12}^{(n)} & [T_{12}^{(n)}, T_{21}^{(1)}] = T_3^{(n)}(\mu) \\ [T_3^{(n)}(\mu), T_{21}^{(1)}] = -2\mu T_{21}^{(n)} & [T_0^{(n)}(\mu), T_{ab}^{(1)}] = 0 \ (n = 1, 2, \ a, b = 1, 2) \end{cases} \tag{2.13}$$

where

$$\begin{cases} T_3^{(n)}(\mu) = T_{22}^{(n)} - \mu T_{11}^{(n)}, & T_0^{(n)}(\mu) = T_{22}^{(n)} + \mu T_{11}^{(n)}, \\ T_+^{(n)} = T_{12}^{(n)}, & T_-^{(n)} = T_{21}^{(n)} \end{cases} \tag{2.14}$$

$$\begin{cases} [T_0^{(2)}(\mu), T_3^{(2)}(\mu)] + 2\mu(T_{21}^{(1)}T_{12}^{(2)} - T_{21}^{(1)}T_{12}^{(1)}) = 0 \\ [T_0^{(n)}(\mu), T_{12}^{(2)}] + T_{12}^{(1)}T_3^{(n)}(\mu) - T_{12}^{(n)}T_3^{(1)}(\mu) = 0 \quad (n \geq 1) \\ [T_0^{(n)}(\mu), T_{21}^{(2)}] + T_{21}^{(n)}T_3^{(1)}(\mu) - T_{21}^{(1)}T_3^{(n)}(\mu) = 0 \quad (n \geq 1) \end{cases} \tag{2.15}$$

and

$$\begin{aligned} T_{12}^{(n+1)} &= (2\mu)^{-1}\{[T_3^{(n)}(\mu), T_{12}^{(2)}] + T_{12}^{(1)}T_0^{(n)}(\mu) - T_{12}^{(n)}T_0^{(1)}(\mu)\} \\ T_{21}^{(n+1)} &= (2\mu)^{-1}\{[T_{21}^{(2)}, T_3^{(n)}(\mu)] + T_{21}^{(1)}T_0^{(n)}(\mu) - T_{12}^{(n)}T_0^{(1)}(\mu)\} \\ T_3^{(n+1)}(\mu) &= [T_{12}^{(n)}, T_{21}^{(2)}] + T_{22}^{(1)}T_{11}^{(n)} - T_{22}^{(n)}T_{11}^{(1)}, \ (n \geq 2) \end{aligned} \tag{2.16}$$

It is emphasized that because of the iterative relation eq. (2.16), only $T^{(1)}$ and $T^{(2)}$ are basic ones. To satisfy all the relations with $T^{(n)}(n \geq 3)$ it is enough to look for the constraints to $T^{(3)}$ that in turn to provide the constraints to $T^{(2)}$ itself. Before doing this let us introduce the "physical" operators:

$$T_{12}^{(1)} = \alpha_+ I_+, \quad T_{21}^{(1)} = \alpha_- I_-, \quad T_3^{(1)}(\mu) = 2\mu I_3, \tag{2.17}$$

$$T_{12}^{(2)} = \beta_+ J_+, \quad T_{21}^{(2)} = \beta_- J_-, \quad T_3^{(2)}(\mu) = 2\beta_3 J_3, \tag{2.18}$$

where $\alpha_\pm$, $\beta_\pm$ and β_3 are constants to be determined.

Substituting eqs. (2.15) and (2.16) into eqs. (2.12) and (2.13) one obtains

$$\alpha_+\alpha_- = \mu, \quad \alpha_+\beta_- = \alpha_-\beta_+ = \beta_3 \tag{2.19}$$

and the algebraic relations

$$\begin{aligned} &[I_3, I_\pm] = \pm I_\pm, \quad [I_+, I_-] = 2I_3 \\ &[I_3, J_\pm] = [J_3, I_\pm] = \pm J_\pm, \quad [I_\pm, J_\mp] = \pm 2J_3 \\ &[I_\pm, J_\pm] = [I_3, J_3] = 0 \end{aligned} \tag{2.20}$$

or by

$$I_\pm = I_1 \pm iI_2, \quad J_\pm = J_1 \pm iJ_2 \tag{2.21}$$

we have

$$\begin{aligned} [I_\lambda, I_\mu] &= C_{\lambda\mu\nu} I_\nu, \quad [I_\lambda, J_\mu] = C_{\lambda\mu\nu} J_\nu, \qquad (2.22)\\ C_{\lambda\mu\nu} &= i\epsilon_{\lambda\mu\nu}, \ (\lambda, \mu, \nu = 1, 2, 3) \qquad (2.23)\end{aligned}$$

Next we have to look for the constraints to $T^{(3)}$ by taking eqs. (2.11)-(2.16) into account. It is easy to know from eq. (2.12) that

$$[T^{(2)}_{22}, T^{(1)}_{11}] = [T^{(3)}_{22}, T^{(2)}_{11}] = 0 \tag{2.24}$$

From eq. (2.16) it follows

$$T^{(3)}_3(\mu) \equiv T^{(3)}_{22} - \mu T^{(3)}_{11} = [T^{(2)}_{12}, T^{(2)}_{21}] + T^{(1)}_{22} T^{(2)}_{11} - T^{(2)}_{22} T^{(1)}_{11} \tag{2.25}$$

By taking the commutator between $T^{(2)}_{22}$ and both of sides of eq. (2.25) and using eq. (2.24) we obtain

$$\mu[T^{(3)}_{11}, T^{(2)}_{22}] = [T^{(2)}_{22}, [T^{(2)}_{12}, T^{(2)}_{21}]] + T^{(1)}_{22}[T^{(2)}_{22}, T^{(2)}_{11}] \tag{2.26}$$

Similarly

$$[T^{(2)}_{11}, T^{(3)}_{22}] = [T^{(2)}_{11}, [T^{(2)}_{12}, T^{(2)}_{21}]] + [T^{(2)}_{22}, T^{(2)}_{11}] T^{(1)}_{11} \tag{2.27}$$

Hence, the relation eq. (2.9) for $[T^{(3)}_{11}, T^{(2)}_{22}] = [T^{(2)}_{11}, T^{(3)}_{22}]$ leads to

$$[T^{(2)}_3(\mu), [T^{(2)}_{12}, T^{(2)}_{21}]] = T^{(1)}_3(\mu)(T^{(1)}_{21} T^{(2)}_{12} - T^{(2)}_{21} T^{(1)}_{12}). \tag{2.28}$$

With the same manipulations to $T^{(3)}_{12}$ in eq. (2.16) one gets

$$\begin{aligned} [T^{(2)}_{12}, T^{(3)}_{12}] &= [T^{(2)}_{12}, [T^{(2)}_{12}, T^{(2)}_{11}]] + T^{(1)}_{12}[T^{(2)}_{12}, T^{(2)}_{11}] - T^{(2)}_{12}[T^{(2)}_{12}, T^{(2)}_{11}] \qquad (2.29)\\ \mu[T^{(3)}_{12}, T^{(2)}_{12}] &= [T^{(2)}_{12}, [T^{(2)}_{22}, T^{(2)}_{12}]] + T^{(1)}_{12}[T^{(2)}_{12}, T^{(2)}_{22}] - T^{(2)}_{12}[T^{(2)}_{12}, T^{(2)}_{22}] \qquad (2.30)\end{aligned}$$

From $[T^{(2)}_{12}, T^{(3)}_{12}] = 0$ it follows

$$[T^{(2)}_{12}, [T^{(2)}_3(\mu), T^{(2)}_{12}]] = T^{(1)}_{12}(T^{(2)}_{12} T^{(1)}_3(\mu) - T^{(1)}_{12} T^{(2)}_3(\mu)) \tag{2.31}$$

Similarly the relations $[T^{(2)}_{21}, T^{(3)}_{21}] = 0$ provides the constraint

$$[T^{(2)}_{21}, [T^{(2)}_3(\mu), T^{(2)}_{21}]] = T^{(1)}_{21}(T^{(2)}_{21} T^{(1)}_3(\mu) - T^{(1)}_{21} T^{(2)}_3(\mu)) \tag{2.32}$$

By taking $[T^{(2)}_{12}, T^{(3)}_3(\mu)]$ and $[T^{(2)}_{21}, T^{(3)}_3(\mu)]$ where $T^{(3)}_3(\mu)$ is given by eq. (2.16) and considering eq. (2.9) we obtain

$$\begin{aligned} &2\mu[T^{(2)}_{12}, [T^{(2)}_{12}, T^{(2)}_{21}]] + [T^{(2)}_3(\mu), [T^{(2)}_3(\mu), T^{(2)}_{12}]]\\ &= (T^{(2)}_{12} T^{(3)}_3(\mu) - T^{(1)}_{12} T^{(2)}_3(\mu))(T^{(1)}_3(\mu) + 2\mu)\\ &\quad -2\mu T^{(1)}_{12}(T^{(2)}_{21} T^{(1)}_{12} - T^{(1)}_{21} T^{(2)}_{12}) \end{aligned} \tag{2.33}$$

and

$$\begin{aligned}&2\mu[T_{21}^{(2)},[T_{12}^{(2)},T_{21}^{(2)}]]+[T_3^{(2)}(\mu),[T_{21}^{(2)},T_3^{(2)}(\mu)]]\\&=(T_{21}^{(2)}T_3^{(3)}(\mu)-T_{21}^{(1)}T_3^{(2)}(\mu))(T_3^{(1)}(\mu)-2\mu)\\&\quad+2\mu T_{21}^{(1)}(T_{21}^{(2)}T_{12}^{(1)}-T_{21}^{(1)}T_{12}^{(2)})\end{aligned}\tag{2.34}$$

we thus have exhausted all the relations including $T^{(3)}$ in eqs. (2.9)-(2.16). Substituting eqs. (2.17) and (2.18) into eqs. (2.28)-(2.34) we derive

$$[J_3,[J_+,J_-]]=\frac{\mu}{\beta_+\beta_-}I_3(J_-I_+-I_-J_+)\tag{2.35}$$

$$[J_\pm,[J_3,J_\pm]]=\frac{\mu}{\beta_+\beta_-}I_\pm(J_\pm I_3-I_\pm J_3)\tag{2.36}$$

$$\begin{aligned}&2[J_3,[J_3,J_\pm]]+[J_\pm,[J_\pm,J_\mp]]\\&=\frac{\mu}{\beta_+\beta_-}\{2I_3(J_\pm I_3-I_\pm J_3)+I_\pm(I_-I_+-J_-I_+)\}\end{aligned}\tag{2.37}$$

By introducing

$$\mu\beta_+^{-1}\beta_-^{-1}=\frac{1}{4}h^2\tag{2.38}$$

the eqs. (2.35)-(2.37) can be recast into

$$\begin{aligned}&[[J_\lambda,J_\mu],[I_\tau,J_\rho]]+[[J_\tau,J_\rho],[I_\lambda,J_\mu]]\\=\;&h^2(a_{\lambda\mu\nu\alpha\beta\gamma}C_{\tau\rho\nu}+a_{\tau\rho\nu\alpha\beta\gamma}C_{\lambda\mu\nu})\{I_\alpha,I_\beta,I_\gamma\}\end{aligned}\tag{2.39}$$

where

$$a_{\lambda\mu\nu\alpha\beta\gamma}=\frac{1}{24}C_{\lambda\rho\sigma}C_{\mu\beta\rho}C_{\nu\gamma\tau}C_{\sigma\gamma\tau}\tag{2.40}$$

$$\{x_1,x_2,x_3\}=\sum_{i\neq j\neq k\neq i}x_ix_jx_k\tag{2.41}$$

With eq. (2.23) it is easy to prove that

$$[J_\lambda,[J_\mu,I_\nu]]-[I_\lambda,[J_\mu,J_\nu]]=0\tag{2.42}$$

In conclusion for given R-matrix eq. (1.15) by solving RTT relation we find the infinite algebra shown by eqs. (2.22), (2.39) and (2.42). This algebra is not closed since J_μ are not closed. Moreover the coproduct Δ is defined by

$$\Delta T_{ab}(u)=\sum_c T_{ac}(u)\otimes T_{cb}(u)\tag{2.43}$$

that by setting $T_{22}^{(1)}=\mu I_3$ and $T_{11}^{(1)}=-\mu I_3$ leads to

$$\Delta(I_\lambda)=I_\lambda\otimes 1+1\otimes I_\lambda\tag{2.44}$$

$$\Delta(J_\lambda)=J_\lambda\otimes 1+1\otimes J_\lambda+\frac{1}{2}hC_{\lambda\mu\nu}I_\nu\otimes I_\mu.\tag{2.45}$$

This set of infinite algebra is called $Y(sl(2))$.

§2.3 Drinfeld's Theory of *Yangian*

Let g be a finite-dimensionally simple Lie algebra the associated algebra A is generated by elements I_λ and J_λ with defining relations

$$[I_\lambda, I_\mu] = C_{\lambda\mu\nu} I_\nu, \quad [I_\lambda, J_\mu] = C_{\lambda\mu\nu} J_\nu, \tag{2.46}$$

$$[J_\lambda, [J_\mu, I_\nu]] - [I_\lambda, [J_\mu, J_\nu]] = h^2 a_{\lambda\mu\nu\alpha\beta\gamma} \{I_\alpha, I_\beta, I_\gamma\} \tag{2.47}$$

$$\begin{aligned}&[[J_\lambda, J_\mu], [I_r, J_s]] + [[J_r, J_s], [I_\lambda, J_\mu]] \\ &= h^2 (a_{\lambda\mu\nu\alpha\beta\gamma} C_{rs\nu} + a_{rs\nu\alpha\beta\gamma} C_{\lambda\mu\nu}) \{I_\alpha, I_\beta, I_\gamma\}\end{aligned} \tag{2.48}$$

where the $C_{\lambda\mu\nu}$ are the structure constants of g, $a_{\lambda\mu\nu\alpha\beta\gamma}$, $\{x_1, x_2, x_3\}$ are given in eqs. (2.40) and (2.41) and all the repeating indices imply summation. Moreover, the co-multiplication is given by

$$\Delta(I_\lambda) = I_\lambda \otimes 1 + 1 \otimes I_\lambda \tag{2.49}$$

$$\Delta(J_\lambda) = J_\lambda \otimes 1 + 1 \otimes J_\lambda + \frac{1}{2} h C_{\lambda\mu\nu} I_\nu \otimes I_\mu \tag{2.50}$$

that are the same as eqs. (2.44) and (2.45) for $sl(2)$. Set eqs. (2.46)-(2.50) is called *Yangian* denoted by $Y(g)$. Setting $h = 1$ it is a Hopf algebra. When $g = sl(2)$ eq. (2.47), i.e., eq. (2.42) is identically zero. However eq. (2.48) holds that has been shown in the above calculation. For $g = sl(n)(n > 2)$ Drinfeld pointed out that [10] eq. (2.48) follows from eqs. (2.46) and (2.47) which is usually called serre relations(There is a mis-print in the remark appeared in p255 of Ref. [10]).

On one word for $sl(2)$ eqs. (2.46) and (2.48) should only be concerned, whereas for $sl(n)(n > 2)$ only eqs. (2.46) and (2.47) should be taken into account. In the §2.1 the $Y(sl(2))$ has explicitly been obtained based on the RTT relation. let us take $Y(sl(2))$ as an example to show the properties of *Yangian*.

1. It contains the classical algebra $sl(2)$ as a sub-algebra denoted by the set $\{I_\lambda\}(\lambda = 1, 2, 3)$ that is closed.

2. Besides the set $\{I_\lambda\}$ there appears a new set $\{J_\lambda\}(\lambda = 1, 2, 3)$ which are not closed. Hence with $\{I_\lambda\}$ and $\{J_\lambda\}$ an infinite algebra can be generated.

3. Observing the expansion eq. (2.3) it looks like a loop algebra with the deference that the $T_{ab}^{(n)}$ obey the "deformed" commutation relations. Thus we may regard *Yangian* as a deformed loop algebra. We have known that both of sides of eq. (2.47) vanish identically for $sl(2)$. The spectral parameter u plays very important role in the *Yangian* theory. Just the spectral parameter dependence gives rise to the new operators $\{J_\lambda\}$ beyond the usual Lie algebra $sl(2)$, and make the infinite algebra not the loop algebra.

4. The RTT relation provides a standard method to generate *Yangian* even though the calculations are tedious. From the RTT relation we have derived the commutation relations satisfied by $\{I_\lambda\}$ and $\{J_\lambda\}$ that are quantum operators. The realizations of such operators can be made through the usual quantum mechanical operators or second quantized operations. The *Yangian* symmetry is model-independent. It is related to a given Lie algebra. However a realization is model-dependent. Hence it is connected with the specified physical model.

§2.4 det$T(u)$ Associated with $Y(gl(2))$

For $sl(2)$ a R-matrix is given by eq. (1.15) and the $T(u)$ is 2 by 2 matrix. Each elements of $T(u)$ matrix is quantum operator. The inverse of $T(u)$ is defined by

$$T^{-1}(u) = \frac{1}{\mathrm{det}T(u)} \begin{bmatrix} \tilde{T}_{11}(u) & \tilde{T}_{12}(u) \\ \tilde{T}_{21}(u) & \tilde{T}_{22}(u) \end{bmatrix} \tag{2.51}$$

and det$T(u)$ should commute with $T_{ab}(u)(a, b = 1, 2)$:

$$[\mathrm{det}T(u), T_{ab}(v)] = 0, \ (a, b = 1, 2), \tag{2.52}$$

i.e., det$T(u)$ is a center of the algebra. From eq. (2.4) by setting $u - v = 1$, it follows

$$[T_{bc}(u), T_{ad}(u-1)] - T_{ac}(u)T_{bd}(u-1) - T_{ac}(u-1)T_{bd}(u) = 0 \tag{2.53}$$

Substituting eq. (2.51) into $T^{-1}(u)T(u) = T(u)T^{-1}(u) = 1$ and making comparison to eq. (2.53) we have

$$\begin{array}{ll} \tilde{T}_{12}(u) = -T_{12}(u-1) & \tilde{T}_{21}(u) = -T_{12}(u-1) \\ \tilde{T}_{11}(u) = T_{22}(u-1) & \tilde{T}_{22}(u) = T_{11}(u-1) \end{array} \tag{2.54}$$

and [10]

$$\mathrm{det}T(u) = T_{11}(u)T_{22}(u-1) - T_{12}(u)T_{21}(u-1). \tag{2.55}$$

The direct check also verifies the validity of eq. (2.52) by virtue of eq. (2.55). Of course, it holds

$$[\mathrm{det}T(u), \mathrm{tr}T(v)] = 0, \ (a, b = 1, 2), \tag{2.56}$$

The conserved family C_n determined by

$$\mathrm{det}T(u) = \sum_{n=0}^{\infty} u^{-n} C_n \tag{2.57}$$

can be calculated explicitly:

$$\begin{aligned} C_0 &= 1, \quad C_1 = T_{11}^{(1)} + \mu T_{22}^{(1)} = T_0^{(1)}(\mu) \\ C_j &= T_0^{(j)}(\mu) + \sum_{\substack{m+l=j \\ m,l\neq 0}} \frac{(l+m-1)!}{(m-1)!l!} T_{22}^{(m)} + \\ & \sum_{\substack{m+n+l=j \\ m,n\neq 0}} \frac{(l+m-1)!}{(m-1)!l!} \left(T_{11}^{(n)} T_{22}^{(m)} - T_{12}^{(n)} T_{21}^{(m)} \right) \ (j \geq 1). \end{aligned} \tag{2.58}$$

The explicit forms of C_k are:

$$C_1 = T_{22}^{(1)} + \mu T_{11}^{(1)} = T_0^{(1)}(\mu) = \mu N, \tag{2.59}$$

$$C_2 = T^{(2)}(\mu) + T_{22}^{(1)} + T_{11}^{(1)} T_{22}^{(1)} - T_{12}^{(1)} T_{21}^{(1)} = \mu(M + \frac{1}{2}N + \frac{1}{2}N^2) \tag{2.60}$$

$$\begin{aligned} C_3 &= T_0^{(3)}(\mu) + T_3^{(2)}(\mu) + T_{11}^{(2)} T_{22}^{(1)} - T_{12}^{(2)} T_{21}^{(1)} + T_{11}^{(1)} T_{22}^{(2)} - T_{12}^{(1)} T_{21}^{(2)} + C_2 \\ &= \frac{1}{2}\mu M N \\ & \cdots \end{aligned} \tag{2.61}$$

where

$$T_0^{(1)}(\mu) \equiv \mu N, \tag{2.62}$$

$$T_0^{(2)}(\mu) \equiv T_{22}^{(2)} + \mu T_{11}^{(2)} = \mu(\vec{I}^2 + M), \tag{2.63}$$

with

$$\vec{I}^2 = \sum_{\lambda=1}^{3} I_\lambda^2 \tag{2.64}$$

the casimir of $gl(2)$ algebra.

It can easily be proved that

$$[C_m, J^{(n)}] = 0, \quad [C_m, C_n] = 0 \tag{2.65}$$

for any μ. Therefore C_m commute with any conserved quantities.

Finally, let us summarize the RTT relation shown by eqs. (2.5) and (2.6) that can be reduced to the following independent set of relations ($T_3^{(n)} = T_3^{(n)}(\mu)$, $T_0^{(n)} = T_0^{(n)}(\mu)$, $T_+^{(n)} = T_{12}^{(n)}$, $T_-^{(n)} = T_{21}^{(n)}$):

$$\begin{cases} [T_\alpha^{(1)}, T_\alpha^{(2)}] = 0, \ (\alpha = \pm, 3) \\ [T_3^{(1)}, T_\pm^{(k)}] = [T_3^{(k)}, T_\pm^{(1)}] = \pm 2\mu T_\pm^{(k)}, \ (k = 1, 2) \\ [T_+^{(1)}, T_-^{(k)}] = [T_+^{(k)}, T_-^{(1)}] = T_3^{(k)}, \end{cases} \tag{2.66}$$

$$\begin{cases} [T_0^{(n)}, T_0^{(m)}] = 0, \ (\forall n, m) \\ [T_0^{(1)}, T_\alpha^{(n)}] = [T_0^{(n)}, T_\alpha^{(1)}] = 0, \ (\alpha = \pm, \ 3), \\ [T_0^{(2)}, T_\pm^{(2)}] = \pm(T_3^{(1)} T_\pm^{(2)} - T_3^{(2)} T_\pm^{(1)}), \\ [T_0^{(2)}, T_3^{(2)}] = 2\mu(T_+^{(1)} T_-^{(2)} - T_+^{(2)} T_-^{(1)}), \end{cases} \tag{2.67}$$

$$\begin{cases} T_\pm^{(n+1)} = (2\mu)^{-1}\{\pm[T_3^{(2)}, T_\pm^{(n)}] + T_0^{(n)} T_\pm^{(1)} - T_0^{(1)} T_\pm^{(n)}\} \\ T_3^{(n+1)} = [T_+^{(n)}, T_-^{(2)}] + (2\mu)^{-1}(T_0^{(n)} T_3^{(1)} - T_0^{(1)} T_3^{(n)}) \end{cases} \tag{2.68}$$

and

$$[T^{(n)}_\alpha, T^{(m)}_\alpha] = 0 \ (\alpha \neq 0,\ m < n,\ n > 2) \tag{2.69}$$

which is equivalent to

$$\begin{cases} [T^{(m)}_\pm, [T^{(2)}_3, T^{(n)}_\pm]] \pm [T^{(m)}_\pm, T^{(n)}_0]T^{(1)}_\pm = 0 \\ 2\mu[T^{(m)}_3, [T^{(n)}_+, T^{(2)}_-]] + [T^{(m)}_3, T^{(n)}_0]T^{(1)}_3 = 0 \end{cases} \tag{2.70}$$

To show the relationship for the set of eqs. (2.66)-(2.69) and the set of eqs. (2.22), (2.42) and (2.39) i.e., the Drinfeld form for $Y(sl(2))$, by taking $m = n = 2$ in eqs. (2.66) and (2.70) it follows

$$\begin{aligned} &2\mu[T^{(2)}_\pm, [T^{(2)}_+, T^{(2)}_-]] \pm [T^{(2)}_3, [T^{(2)}_3, T^{(2)}_\pm]] \\ &= 2\mu(T^{(1)}_+ T^{(2)}_- - T^{(2)}_+ T^{(1)}_-)T^{(1)}_\pm \pm (T^{(1)}_3 T^{(2)}_\pm - T^{(2)}_3 T^{(1)}_\pm)T^{(1)}_3 \end{aligned} \tag{2.71}$$

Eqs. (2.66), (2.31) and (2.34) lead to eqs. (2.22), (2.39) and (2.42), i.e., $Y(sl(2))$ can be generated based on $T^{(1)}$ and $T^{(2)}$. The "closure" of eqs. (2.22), (2.39) and (2.42) for $sl(2)$ means that we can use $\{I_\lambda\}$ and $\{J_\lambda\}$ to generate any new generators of $Y(sl(2))$ based on eq. (2.68) to form an infinite algebra.

§3 Realization of *Yangian*

§3.1 $Y(sl(2))$

$$\vec{I} = \sum_{i=1}^{N} \vec{S}_i, \quad \vec{J} = \sum_{i\neq j}^{N} \vec{S}_i \times \vec{S}_j \tag{3.1}$$

where $\vec{I} = \{I_\lambda\}$ and $\vec{J} = \{J_\lambda\}(\lambda = 1, 2, 3)$ and

$$[S^\lambda_i, S^\mu_j] = i\epsilon_{\lambda\mu\nu} S^\nu_i \delta_{ij} \tag{3.2}$$

satisfy eqs. (2.46)-(2.48). Therefore eq. (3.1) forms a $Y(sl(2))$ [11].

To prove the statement first we rewrite eq. (3.2)

$$\begin{aligned} &[S^\alpha_i, S^\alpha_j] = 0, \quad \text{for } \alpha = \pm,\ 3, \\ &[S^3_i, S^\pm_j] = \pm\delta_{ij} S^\pm_i, \quad [S^+_i, S^-_j] = 2\delta_{ij} S^3_i \end{aligned}$$

We should verify that the defined $\{I_\lambda\}$ and $\{J_\lambda\}$ satisfy eq. (2.20) and eqs. (2.35)-(2.37). Since by $J_\lambda \rightarrow h^{\frac{1}{2}} J_\lambda$ the constant h can be removed, one can takes $h = 1$.

Defining

$$\begin{aligned} &I_\pm = \sum_i S^\pm_i, \qquad I_3 = \sum_i S^3_i, \\ &J_\pm = \mp \sum_{i\neq j} W_{ij} S^\pm_i S^3_j, \quad J_3 = \frac{1}{2}\sum_{i\neq j} W_{ij} S^+_i S^-_j \end{aligned} \tag{3.3}$$

where $W_{ij} + W_{ji} = 0$, and substituting them into eqs. (2.20), (2.35)-(2.37) we have

$$(S_i^{\pm})^2 = 0, \quad (S_i^3)^2 = 1$$

i.e., the operators S_i^{λ} should be Pauly matrices, and

$$W_{ij}W_{jk} + W_{jk}W_{ki} + W_{ki}W_{ij} = -h^2, \text{ (for } i \neq j \neq k \neq i)] \tag{3.4}$$

The simplest solution of W_{ij} is given by

$$W_{ij} = \begin{cases} h & i < j \\ 0 & i = j \\ -h & i > j \end{cases} \tag{3.5}$$

that is nothing but the statement.

§3.2 Example for $Y(sl(2))$

By using the fermion operators acting on i-th site

$$a_i^{\dagger} = c_{i\uparrow}^{\dagger}, \quad a_i = c_{i\uparrow}, \quad b_i^{\dagger} = c_{i\downarrow}^{\dagger}, \quad b_i = c_{i\downarrow} \tag{3.6}$$

satisfying

$$\{a_i, a_j\} = \{a^{\dagger}, a_j^{\dagger}\} = \{b_i, b_j\} = \{b_i^{\dagger}, b_j^{\dagger}\} = 0 \tag{3.7}$$

$$\{a_i^{\dagger}, a_j\} = \delta_{ij}, \quad \{b_i^{\dagger}, b_j\} = \delta_{ij}, \tag{3.8}$$

and $a_i^{\dagger}(a_i)$ anticommute with $b_i^{\dagger}(b_i)$. The realization of $Y(sl(2))$ can be made by

$$\begin{aligned} I_+ &= \sum_i a_i^{\dagger} b_i = \sum_i I_i^+ \\ I_- &= \sum_i b_i^{\dagger} a_i = \sum_i I_i^- \\ I_3 &= \frac{1}{2}\sum_i (a_i^{\dagger} a_i - b_i^{\dagger} b_i) = \sum_i I_i^3 \end{aligned} \tag{3.9}$$

that satisfy $sl(2)$ algebra

$$[I_3, I_{\pm}] = \pm I_{\pm}, \quad [I_+, I_-] = 2I_3 \tag{3.10}$$

and

$$\begin{aligned} J_+ &= \sum_{i,j} \theta_{i,j} a_i^{\dagger} b_j - U \sum_{i,j} \epsilon_{i,j} I_i^+ I_j^3 = \sum_i J_i^+ \\ J_- &= \sum_{i,j} \theta_{i,j} b_i^{\dagger} a_j + \sum_{i,j} \epsilon_{i,j} I_i^- I_j^3 = \sum_i J_i^- \\ J_3 &= \frac{1}{2}\left\{ \sum_{i,j} \theta_{i,j} (a_i^{\dagger} a_j - b_i^{\dagger} b_j) + U \sum_{i,j} \epsilon_{i,j} I_i^+ I_j^- \right\} = \sum_i J_i^3 \end{aligned} \tag{3.11}$$

where

$$\theta_{i,j} = \delta_{i,j-1} - \delta_{i,j+1}, \quad \epsilon_{ij} = \begin{cases} 1 & i < j \\ 0 & i = j \\ -1 & i > j \end{cases} \tag{3.12}$$

By virtue of eqs. (3.6)-(3.8) the validity of eq. (2.22) for eqs. (3.10)-(3.12) is convinced. One then able to set

$$T_{\pm}^{(1)} = \alpha_{\pm} I_{\pm}, \ T_3^{(1)} = 2\alpha_+\alpha_- I_3, \ T_{\pm}^{(2)} = \beta_{\pm} J_{\pm}, \ T_3^{(2)} = 2\alpha_+\beta_- J_3 \tag{3.13}$$

with

$$\alpha_+\alpha_- = \mu, \quad \beta_+\alpha_- = \alpha_+\beta_-. \tag{3.14}$$

We thus have made a physical realization of eq. (2.66). Since the construction of $T_0^{(n)}(\mu)$ has been made the eq. (2.67) hold, we should show the validity of eq. (2.70) for eqs. (3.9)-(3.12). Substituting eqs. (3.13) and (3.14) into eq. (2.70) and calculating order by order we find the forms of the formulae of Drinfeld for $m = n = 2$:

$$\begin{aligned} [J_{\pm}, [J_3, J_{\pm}]] &= \frac{\mu}{\beta_+\beta_-}(I_3 J_{\pm} - J_3 I_{\pm}) I_{\pm}, \\ [J_3, [J_+, J_-]] &= \frac{\mu}{\beta_+\beta_-}(I_+ J_- - J_+ I_-) I_3 \end{aligned}$$

where

$$\mu \beta_+^{-1} \beta_-^{-1} = U^2.$$

For $m = n + 1$ and $m = n - 1$ one can check that eq. (2.69) identically vanishes. It means that eq. (2.69) is satisfied for $m = 3, \ n = 2$ and $m = 2, \ n = 3$. By direct calculation eq. (2.69) can be shown to be true for $m = 2, \ n = 4$ and $m = n = 3$.

What is the physical meaning of such a realization of $\{I_\lambda\}$ and $\{I_\lambda\}(\lambda = 1, 2, 3)$? The $\{\vec{I_i}\}$ describe the local spin "rotation"(up-down) and do not shift the fermion located at lattice. However the operators $\vec{J_i}$ describe the hopping of fermion from the i-th site to the next ones. More explicitly [12] eq. (3.11) can be written as

$$\begin{aligned} J_+ &= \sum_{i,j}(a_i^\dagger b_{i+1} - a_i^\dagger b_{i-1}) - U \sum_{i,j}\{a_i^\dagger b_i (a_j^\dagger a_j - b_j^\dagger b_j) - a_j^\dagger b_j (a_i^\dagger a_i - b_i^\dagger b_i)\}, \\ J_3 &= \frac{1}{2}\sum_{i,j}(a_i^\dagger a_{i+1} - b_i^\dagger b_{i+1} - a_i^\dagger a_{i-1} + b_i^\dagger b_{i-1}) + U \sum_{i,j} a_i^\dagger b_i b_j^\dagger a_j, \\ J_- &= \sum_{i,j}(b_i^\dagger a_{i+1} - b_i^\dagger a_{i-1}) + U \sum_{i,j}\{b_i^\dagger a_i (a_j^\dagger a_j - b_j^\dagger b_j) - b_j^\dagger a_j (a_i^\dagger a_i - b_i^\dagger b_i)\}. \end{aligned} \tag{3.15}$$

The realization eq. (3.15) possesses a nice property in that the Hamiltonian of the 1-dim. Hubbard model

$$H = \sum_i (a_i^\dagger a_{i+1} + a_{i+1}^\dagger a_i + b_i^\dagger b_{i+1} + b_{i+1}^\dagger b_i) - U \sum_i (a_i^\dagger a_i - \frac{1}{2})(b_i^\dagger b_i - \frac{1}{2}) \tag{3.16}$$

commutes with $\{I_\lambda\}$ and $\{J_\lambda\}$ given by eqs. (3.9) and (3.15):

$$\begin{aligned} [H, I_\alpha] &= 0, && (3.17)\\ [H, J_\alpha] &= 0, \ (\alpha = \pm, 3) && (3.18) \end{aligned}$$

Therefore

$$[H, Y(sl(2))] = 0 \tag{3.19}$$

i.e., $Y(sl(2))$ serves as a symmetry of 1-d Hubbard model. The eq. (3.17) is easy to be proved. For eq. (3.18) the periodic boundary condition should be taken account. By substituting eqs. (3.15) and (3.16) into the commutator eq. (3.18) the direct check verifies the validity of the statement [12].

For a nonlinear model with hopping there exists the interaction between different sites. Is there a global symmetry besides the "total" spin? The answer is yes. This new symmetry is nothing but the $Y(sl(2))$ in the example. To form a *Yangian* the operators $\{J_\lambda\}$ play the central role. They describe the hopping from i-th site to $(i+1)$ or $(i-1)$ plus the operators $\sum_{i\neq j}\vec{S}_i \times \vec{S}_j$. Obviously $\{J_\lambda\}$ cannot be closed, because the set $\{J_\lambda\}$ shifts the fermion one by one site. Only at the periodic boundary it makes the "particle" return to the starting one. The existence of *Yangian* means that we can move a "particle" to any site through only two types of operations: "local rotation"($\{I_\lambda\}$) and "one-site shift" ($\{J_\lambda\}$). The consistence of associativity for the successive shift operation yields the constraint condition to $\{I_\lambda\}$ and $\{J_\lambda\}$, as shown by eq. (2.39). For a physical system if there is $sl(2)$(or $SU(2)$) symmetry in the absence of interaction, we may ask whether there exists symmetries that describe the nonlinear-interaction systems. For the particular classes of models the answer is yes. The *Yangian* is just served as such a symmetry.

§3.3 Example for $Y(sl(n))(n > 2)$

Let us discuss an example for long-ranged interaction model associated with $sl(n)(n > 2)$ [13]. Introducing the spin operators

$$I_a = \sum_{i=1}^{N} S_i^a, \ (a = 1, \ldots, n) \tag{3.20}$$

where

$$[S_i^a, S_j^b] = f_{abc} S_i^c \delta_{ij} \tag{3.21}$$

and

$$J_a = \sum_{i\neq j} W_{ij} f_{abc} S_i^b S_j^c, \ (W_{ij} = -W_{ji}) \tag{3.22}$$

The structure constants f_{abc} obey the Jacobi identity:

$$f_{abc}f_{dec} = f_{adc}f_{bec} - f_{aec}f_{bdc}. \tag{3.23}$$

It is easy to verify that eq. (2.46) is satisfied with $C_{abc} = f_{abc}$. As pointed by Drinfeld for $sl(n)(n > 2)$ eqs. (2.46) and (2.47) imply eq. (2.48). Therefore we should check the validity of eq. (2.47) based on eqs. (3.20) and (3.22). After calculation we get

$$[J_a, J_b] = \sum_{i\neq j\neq k\neq i}^{N} W_{ij}W_{ik}f_{\alpha\beta\mu}f_{b\gamma\nu}f_{\alpha\mu\nu}T^{\alpha\beta\gamma}_{ijk} \tag{3.24}$$

where

$$T^{\alpha\beta\gamma}_{ijk} = S^\beta_j S^\alpha_i S^\gamma_k + S^\beta_j S^\gamma_k S^\alpha_i + S^\gamma_k S^\alpha_i S^\beta_j + S^\alpha_i S^\gamma_k S^\beta_j \tag{3.25}$$

Hence

$$[J_a, [J_b, J_c]] = f_{bc\lambda}[J_a, J_\lambda] = \sum_{i\neq j\neq k\neq i}^{N} W_{ij}W_{ik}g^{\alpha\beta\gamma}_{abc}T^{\alpha\beta\gamma}_{ijk} \tag{3.26}$$

where

$$g^{\alpha\beta\gamma}_{abc} = f_{\alpha\mu\nu}f_{a\beta\mu}f_{bc\lambda}f_{\lambda\gamma\nu} \tag{3.27}$$

repeated indices are summed. One can prove that

$$(g^{\alpha\beta\gamma}_{abc} + g^{\alpha\beta\gamma}_{bca} + g^{\alpha\beta\gamma}_{cab}) + (g^{\alpha\gamma\beta}_{abc} + g^{\alpha\gamma\beta}_{bca} + g^{\alpha\gamma\beta}_{cab}) = \sum t^{(\alpha\beta\gamma)}_{abc} \tag{3.28}$$

where $(\alpha\beta\gamma)$ denotes the summation over the permutation to $\alpha,\ \beta$ and γ

$$t^{\alpha\beta\gamma}_{abc} = f_{a\alpha\lambda}f_{b\beta\mu}f_{c\gamma\nu}f_{\lambda\mu\nu} = 4!\ a_{abc\alpha\beta\gamma} \tag{3.29}$$

that has appeared in eq. (2.40).

With the notations we have

$$\sum_{i\neq j\neq k\neq i}^{N} W_{ij}W_{ik}(g^{\alpha\beta\gamma}_{abc} + g^{\alpha\beta\gamma}_{bca} + g^{\alpha\beta\gamma}_{cab})T^{\alpha\beta\gamma}_{ijk} = \sum_{i\neq j\neq k\neq i}^{N} W_{ij}W_{ik}t^{(\alpha\beta\gamma)}_{abc}T^{\alpha\beta\gamma}_{ijk} \tag{3.30}$$

With the help of the above relations on the basis of the Jacobi identities

$$[J_a, [J_b, I_c]] + [I_a, [J_b, J_c]] = [J_a, [J_b, I_c]] + [J_c, [I_a, J_b]] + [J_b, [J_c, I_a]]. \tag{3.31}$$

Only the case where $a \neq b \neq c \neq a$ should be taken into account. These components of eq. (3.31) are equal to

$$\begin{aligned} 2\ \ {\textstyle\sum_{i\neq j\neq k}^{N}} \quad & (\ W_{ij}W_{ik}t^{\alpha\beta\gamma}_{abc}S^\alpha_i S^\beta_j S^\gamma_k + W_{ik}W_{ij}t^{\alpha\beta\gamma}_{abc}S^\alpha_i S^\gamma_k S^\beta_j \\ & + W_{ki}W_{kj}t^{\alpha\beta\gamma}_{abc}S^\gamma_k S^\alpha_i S^\beta_j + W_{ji}W_{jk}t^{\alpha\beta\gamma}_{abc}S^\beta_j S^\alpha_i S^\gamma_k \\ & + W_{jk}W_{ji}t^{\alpha\beta\gamma}_{abc}S^\beta_j S^\gamma_k S^\alpha_i + W_{kj}W_{ki}t^{\alpha\beta\gamma}_{abc}S^\gamma_k S^\beta_j S^\alpha_i) \\ = \quad & \frac{2}{3}\sum_{i\neq j\neq k\neq i}^{N}(W_{ij}W_{ik} + W_{jk}W_{ji} + W_{ki}W_{kj})t^{\alpha\beta\gamma}_{abc}\{S^\alpha_i, S^\beta_j, S^\gamma_k\} \end{aligned} \tag{3.32}$$

Defining

$$\Delta_{ijk} = W_{ij}W_{ik} + W_{jk}W_{ji} + W_{ki}W_{kj},\ (W_{ij} = -W_{ji}) \tag{3.33}$$

and noting that there is no summation over the repeated lattice indices, when

$$\Delta_{ijk} = h^2,\quad \text{(independent of } i, j, k) \tag{3.34}$$

the lhs of eq. (3.31)(with $a \neq b \neq c \neq a$) is equal to

$$h^2 a_{abcdef}\{I^d, I^e, I^f\} \tag{3.35}$$

which is exactly the relation eq. (2.47). Thus let eq. (3.22) satisfy $Y(sl(2))$ the eq. (3.33) should be satisfied, i.e.,

$$\Delta_{ijk} = W_{ij}W_{ik} + W_{jk}W_{ji} + W_{ki}W_{kj} = h^2,\ (\forall\ i \neq j \neq k \neq i) \tag{3.36}$$

which is the same as eq. (3.4). Besides the solution (3.5), the general solution [14]

$$W_{jk} = (ih)\frac{Z_j + Z_k}{Z_j - Z_k} \tag{3.37}$$

where $\{Z_j\}$ is a discrete set of complex number. Thus

$$J_a = U\sum_{i=1}^{N} S_i^a + ih\sum_{j\neq k}^{N} \frac{Z_j + Z_k}{Z_j - Z_k} f_{abc}S_j^b S_k^c,\ (a, b, c = 1, 2, \ldots, n) \tag{3.38}$$

together with eq. (3.20) generate the $Y(sl(n))$ for a free parameter U. Here we have used the important property of *Yangian* [10, 14]:

Suppose $\{I_a\}$ and $\{J_a\}$ generate *Yangian* then the $\{I_a\}$ and $\{\tilde{J}_a\}$ also generate a *Yangian*, where

$$\tilde{J}_a = UI_a + J_a, \tag{3.39}$$

U being an arbitrary parameter. This statement can be verified easily because $\tilde{J}_a$ behaves completely in the same manner as J_a.

The simplest form of W_{jk} is $Z_k = e^{i\frac{k\pi}{N}}$ (k integer), then

$$W_{jk} = h\coth\frac{(j-k)\pi}{N}. \tag{3.40}$$

The associated Hamiltonian systems were found in Refs. [15]. Now we can deduce the Hamiltonian from our point of view.

Noting that

$$C_2 = T_0^{(2)} - \frac{1}{2}(2I_3^2 + I_+I_- + I_-I_+) + \frac{1}{2}[T^{(1)} + \frac{1}{2}(T_0^{(1)})^2] \tag{3.41}$$

Assume

$$T_0^{(2)} = \sum_{i\neq j} f_{ij}(P_{ij} - 1) \tag{3.42}$$

where P_{ij} stand for the permutation operators

$$P_{ij} = 2S_i^3 S_j^3 + S_i^+ S_j^- + S_i^- S_j^+ - \frac{1}{2} \tag{3.43}$$

then from

$$\begin{aligned} [T_0^{(2)}, T_\pm^{(2)}] &= \pm[T_3^{(1)} T_\pm^{(2)} - T_3^{(2)} T_\pm^{(1)}] \\ [T_0^{(2)}, T_3^{(2)}] &= 2[T_+^{(1)} T_-^{(2)} - T_+^{(2)} T_-^{(1)}] \end{aligned}$$

it follows

$$f_{ij} = \frac{1}{2} W_{ij}^2$$

we obtain

$$C_2 = -2H_2 + N(N-1) + \frac{1}{2}[T_0^{(1)} + \frac{1}{2}(T_0^{(1)})^2] \tag{3.44}$$

where

$$H_2 = -\frac{1}{4} \sum_{i \neq j} (W_{ij}^2 - 1)(P_{ij} - 1) \tag{3.45}$$

and $T_0^{(1)},\ C_2,\ \{I_\alpha\}$ and $\{J_\alpha\}(\alpha = \pm,\ 3)$ are all commutative. Therefore we have the Hamiltonian

$$H_2 = \sum_{i \neq j} \frac{Z_i Z_j}{Z_{ij} Z_{ji}} (P_{ij} - 1),\ (Z_{ij} = Z_i - Z - j) \tag{3.46}$$

that was obtained in Ref. [15]

§4 Long-Range Interaction Models and *Yangian*

§4.1 Polychronakos' Approach

Besides the nearest neighborhood models(such as Heisenberg chain, Hubbard model, ...), recently, a number of one-dimensional long-range interaction models have been studied [15]. The typical one is Calogero-Sutherland model [16], then it is subsequently extended to the models with internal spin degrees of freedom [17]. Among them an interesting approach was proposed by Bernard, Gaudin, Haldane and Pasquier (BGHP)[17] who made this type of models related to the RTT relation associated with Yang-Baxter equation(YBE). The BGHP approach provides a method to deal with long-range interaction models: for a given rational solution of YBE, for example, $R(u) = u + P$ which is given in eq. (1.15), RTT relation gives rise to the *Yangian* symmetry. With a particular realization of the *Yangian*, in general, we can generate corresponding Hamiltonian of the considered systems.

On the other hand Polychronakos had formulated the integrability in terms of the "coupled" momentum operators [18]:

$$\pi_i = p_i + i\sum_{j\neq i} V_{ij}K_{ij} \tag{4.1}$$

where $p_i = -i\frac{\partial}{\partial x_i}$ ($\hbar = 1$), $V_{ij} = V(x_i - x_j)$ a potential to be determined and K_{ij} the particle permutation operators. The requirements of the Hermiticity of π_i, the absence of linear terms in p_i and that only the two-body potentials in the Hamiltonian lead to

$$\begin{aligned} &V(x) = -V(-x)\ , \\ &H_0 \equiv \frac{1}{2}\sum_i \pi_i^2 = \sum_i p_i^2 + \frac{1}{2}\sum_{i\neq j}\left[\frac{\partial}{\partial x_i}V_{ij}K_{ij} + V_{ij}^2\right] - \frac{1}{6}\sum_{i\neq j\neq k\neq i} V_{ijk}K_{ijk} \end{aligned} \tag{4.2}$$

where

$$\begin{aligned} V_{ijk} &= V_{ij}V_{jk} + V_{jk}V_{ki} + V_{ki}V_{ij} = W_{ij} + W_{jk} + W_{ki}\ , \\ K_{ijk} &= K_{ij}K_{jk} \end{aligned} \tag{4.3}$$

with $W_{ij} = W(x_i - x_j)$ being a symmetric function. The commutation relation between π_i and π_j is found to be

$$[\pi_i, \pi_j] = \sum_{k\neq i,j} V_{ijk}(K_{ijk} - K_{jik})\ . \tag{4.4}$$

This approach can be applied to many integrable systems, including the CS model, δ-interaction model and so on.

§4.2 Sutherland-Römer and Yan Models

Let us first discuss the extended forms of V_{ij} in eq. (4.1) that are different from those given by Ref. [18]. Setting

$$V_{ij} = P_{ij}^{+}a_{ij} + P_{ij}^{-}b_{ij} \tag{4.5}$$

where $P_{ij}^{\pm}$ are given as

$$P_{ij}^{\pm} = \frac{1 \pm \sigma_i\sigma_j}{2}\ (\sigma_i^2 = 1) \tag{4.6}$$

σ_i is quantum operators obeying

$$\begin{aligned} \sigma_i K_{ij} &= K_{ij}\sigma_j\ , \\ \sigma_i K_{mn} &= K_{mn}\sigma_i\ \ (i \neq m, n)\ , \end{aligned}$$

then by substituting eq. (4.5) into eq. (4.1) and doing the parallel discussion in Ref. [18], we find

$$V_{ijk} = P_{ijk}^{+}A_{ijk} + P_{ijk}^{-}A_{ijk} + P_{kij}^{-}A_{kij} + P_{jki}^{-}A_{jki} \tag{4.7}$$

where

$$\begin{aligned} P^{\pm}_{ijk} &= P^{\pm}_{ij} P^{\pm}_{ik} \ , \\ A_{ijk} &= a_{ij}a_{jk} + a_{jk}a_{ki} + a_{ki}a_{ij} \ , \\ B_{ijk} &= a_{ij}b_{jk} + b_{jk}a_{ki} + b_{ki}a_{ij} \ . \end{aligned}$$

Noting that $P^+_{ijk} = P^+_{ikj} = \cdots = P^+_{kji}$, but $P^-_{ijk} = P^-_{jik}$ only.

The sufficient condition of the quantum integrability of eq. (4.1) is

$$V_{ijk} = \text{constant} \quad (\text{or} \quad \text{zero}) \ . \tag{4.8}$$

Now let us look for new solution of eq. (4.8)

(1) When $A_{ijk} \neq 0$, $B_{ijk} \neq 0$, a sufficient solution can be checked:

$$\begin{aligned} &a(x) = l \coth(ax) \qquad (\text{or } a(x) = l \cot(ax)) \ , \\ &b(x) = l \tanh(ax) \qquad (\text{or } b(x) = l \tan(ax)) \end{aligned} \tag{4.9}$$

where $x \equiv x_{ij} = x_i - x_j$, $a, \ l$ are constants and

$$V_{ijk} = -l^2 (P^+_{ijk} + P^-_{ijk} + P^-_{kij} + P^-_{jki}) = -l^2 \ . \tag{4.10}$$

Define

$$H = \frac{1}{2} \sum_i \pi_i^2 - \frac{l^2}{6} \sum_{i \neq j \neq k \neq i} K_{ijk} \ , \tag{4.11}$$

then eq. (4.10) leads to

$$H = \frac{1}{2} \sum_i p_i^2 + \sum_{i<j} l(l - aK_{ij}) \left[\frac{P^+_{ij}}{\sinh^2(ax_{ij})} - \frac{P^-_{ij}}{\cosh^2(ax_{ij})} \right] \ . \tag{4.12}$$

When $K_{ij} = \pm 1$, eq. (4.12) was firstly given in Ref. [19] and studied by Sutherland and Römer [20] in detail.

Define

$$\bar{\pi}_i = \pi_i + il \sum_{i \neq j} K_{ij} \ , \tag{4.13}$$

then

$$[\bar{\pi}_i, \bar{\pi}_j] = 2il(\bar{\pi}_i - \bar{\pi}_j) K_{ij} \tag{4.14}$$

$$[H, \pi_i] = [H, \bar{\pi}_i] = 0 \ . \tag{4.15}$$

The conserved quantities are given by

$$I_n = \sum_i \bar{\pi}_i^n \tag{4.16}$$

which leads to

$$\begin{aligned} [I_n, I_m] &= 0 \ , & (4.17) \\ [H, I_n] &= 0 \ , & (4.18) \end{aligned}$$

i.e. the model is quantum integrable in the sense of Liouville.

(2) When $B_{ijk} = 0$, we consider two cases

(a) $A_{ijk} = 0$

$$\begin{aligned} a(x) &= \frac{l}{x} \ , \quad V_{ijk} = 0 \ , \\ H &= \frac{1}{2}\sum_i p_i^2 + \frac{1}{2}\sum_{i\neq j} \frac{l(l-K_{ij})}{(x_i - x_j)^2} P_{ij}^+ \qquad (4.19) \end{aligned}$$

that is well known as Calogero model when P_{ij}^+ takes the value 1.

(b) $A_{ijk} = \beta^2 \neq 0$

$$[\pi_i, \pi_j] = \beta \sum_{k\neq i,i} P_{ijk}^+ (K_{ijk} - K_{jik}) \ . \qquad (4.20)$$

Define

$$\bar{\pi}_i = \pi_i + \beta \sum_{i\neq j} P_{ij}^+ K_{ij} \ , \qquad (4.21)$$

it is easy to prove that

$$[\bar{\pi}_i, P_{jk}^+] = 0, \quad \forall i \text{ and } j \neq k \qquad (4.22)$$

and

$$\begin{aligned} [\bar{\pi}_i, \bar{\pi}_j] &= 2\beta P_{ij}^+ (\bar{\pi}_i - \bar{\pi}_j) K_{ij} \ , & (4.23) \\ [\bar{\pi}_i^n, \bar{\pi}_j] &= 2\beta P_{ij}^+ (\bar{\pi}_i^n - \bar{\pi}_j^n) K_{ij} \ , & (4.24) \end{aligned}$$

so that eq. (4.17) is also satisfied. Let

$$H = \frac{1}{2}\sum_i \pi_i^2 + \frac{\beta^2}{6} \sum_{i\neq j\neq k\neq i} P_{ijk}^+ K_{ijk} \ . \qquad (4.25)$$

With the help of eq. (4.20), one can prove

$$[H, \pi_i] = [H, \bar{\pi}_i] = [H, I_n] = 0 \ . \qquad (4.26)$$

For the case (b) we have two sufficient solutions of V_{ijk}:

(i) $a(x) = il\cot(ax)$ (or$a(x) = l\coth(ax)$,

$$\begin{aligned} &V_{ijk} = -l^2 P_{ijk}^+ \ , \\ &H = \frac{1}{2}\sum_i p_i^2 + \frac{1}{2}\sum_{i\neq j} \frac{l(l-aK_{ij})}{\sin^2 a(x_i - x_j)} P_{ij}^+ \ . \qquad (4.27) \end{aligned}$$

Eq. (4.27) is the generalization of the spin chain model considered by BGHP [17].
(ii) $a(x) = l\mathrm{sgn}(x)$,

$$H = \frac{1}{2}\sum_i p_i^2 + \frac{1}{2}\sum_{i\neq j} l(l-K_{ij})\delta(x_i-x_j)P_{ij}^+ \ . \tag{4.28}$$

with the condition that $K_{ij} = \pm 1$, eq. (4.28) was first pointed out by Yan [21] through Bethe Ansatz, he also found the Y-operator defined by Yang [6] for eq. (4.28)

$$Y_{ij}^{\alpha\beta} = \frac{1}{ik_{ij}(ik_{ij}-2c)}[ik_{ij} - c(1-\sigma_i\sigma_j)][-ik_{ij}P^{\alpha\beta} + c(1+\sigma_i\sigma_j)] \tag{4.29}$$

where P is the permutation and Y satisfies

$$Y_{jk}^{\alpha\beta}Y_{ik}^{\beta\gamma}Y_{ij}^{\alpha\beta} = Y_{ij}^{\beta\gamma}Y_{ik}^{\alpha\beta}Y_{jk}^{\beta\gamma} \tag{4.30}$$

and $c = l(l\pm 1)/2$ for $K_{ij} = \pm 1$. Noting that there is only P_{ij}^+ in the Hamiltonian eq. (4.28) for the quantum integrability.

Now we have re-interpreted the models eq. (4.12) and eq. (4.28) from the point of view of the formulation eq. (4.1). Next we shall set up the *Yangian* description of models eq. (4.12) and eq. (4.28) through RTT relation.

§4.3 RTT Relation and Long-Range Interaction Models

Let us apply the BGHP approach [17] to the S-R model and Yan model. The solution of Yang-Baxter equation, R-matrix, takes the simplest form (1.15) as

$$R(u) = u + \lambda P_{00'} \tag{4.31}$$

and the RTT relation reads

$$R_{00'}(u-v)T^0(u)T^{0'}(v) = T^{0'}(v)T^0(u)R_{00'}(u-v) \tag{4.32}$$

where $T^0(u) = T(u)\otimes 1$, $T^{0'} = 1\otimes T(u)$ and $P_{00'}$ is the permutation operator exchanging the two auxiliary spaces 0 and $0'$. Make the expansion

$$\begin{aligned} T^0(u) &= I + \sum_{a,b=1}^{p} X_{ba}^0 \sum_{n=0}^{\infty}\frac{\lambda T_n^{ab}}{u^{n+1}} \ , & (4.33) \\ P_{00'} &= \sum_{a,b=1}^{p} X_{ba}^0 X_{ab}^{0'} \ . & (4.34) \end{aligned}$$

Notice that T_n^{ab} is the same as $T_{ba}^{(n+1)}$ in the §1 and §2. It is well known that $\{T_n^{ab}\}$ generate the *Yangian* [10]. Substituting eqs. (4.31), (4.33) and (4.34) into eq. (4.32) one finds

$$\sum_{a,b}\sum_{c,d} X_{ba}^0 X_{dc}^{0'} \sum_{n=0}^{\infty}\left\{u^{-n-1}f_1^n - v^{-n-1}f_2^n + \sum_{m=0}^{\infty} u^{-n-1}v^{-m-1}f_3^{n,m}\right\} = 0 \tag{4.35}$$

where

$$\begin{aligned} f_1^n &= \delta_{bc}T_n^{ad} - \delta_{ad}T_n^{cb} - [T_n^{ab}, T_0^{cd}] \ , \\ f_2^n &= \delta_{bc}T_n^{ad} - \delta_{ad}T_n^{cb} - [T_0^{ab}, T_n^{cd}] \ , \\ f_3^{n,m} &= \lambda(T_n^{ad}T_m^{cb} - T_m^{ad}T_n^{cb}) + [T_{n+1}^{ab}, T_m^{cd}] - [T_n^{ab}, T_{m+1}^{cd}] \ . \end{aligned}$$

For any auxiliary space $\{X_{ab}\}$ we require $f_1^n = f_2^n = f_3^{n,m} = 0$. Obviously, $f_1^n = 0$ is equivalent to $f_2^n = 0$. So we need only to take

$$f_1^n = f_3^{n,m} = 0 \tag{4.36}$$

into account.

First from $f_3^{n,0} = 0$ it follows

$$\delta_{bc}T_{n+1}^{ad} - \delta_{ad}T_{n+1}^{cb} = \lambda(T_0^{ad}T_n^{cb} - T_n^{ad}T_0^{cb}) + [T_n^{ab}, T_1^{cd}] \tag{4.37}$$

which can be recast to

$$T_{n+1}^{ad} = \lambda(T_0^{ad}T_n^{cc} - T_n^{ad}T_0^{cc}) + [T_n^{ac}, T_1^{cd}] \ (a \neq d) \ , \tag{4.38}$$

$$T_{n+1}^{aa} - T_{n+1}^{cc} = \lambda(T_0^{aa}T_n^{cc} - T_n^{aa}T_0^{cc}) + [T_n^{ac}, T_1^{ca}] \ , \tag{4.39}$$

where no summation for the repeating indices is taken. Eqs. (4.38) and (4.39) imply that T_n^{ab} can be determined by iteration for given T_0^{ab} and T_1^{ab}.

Now let us set

$$T_0^{ab} = \sum_{i=1}^{N} I_i^{ab} \ , \tag{4.40}$$

$$T_1^{ab} = \sum_{i=1}^{N} I_i^{ab} D_i \tag{4.41}$$

and

$$[I_i^{ab}, I_j^{cd}] = \delta_{ij}(\delta_{bc}I_i^{ad} - \delta_{ad}I_i^{cb}) \tag{4.42}$$

where D_i are operators to be determined. Substituting eqs. (4.40)–(4.42) into f_1^1 we obtain

$$\sum_i \sum_j I_i^{ab}[D^i, I_j^{cd}] = 0 \ . \tag{4.43}$$

Further we assume

$$\sum_i I_i^{ab}[D_i, I_j^{cd}] = 0, \text{for any } j \tag{4.44}$$

with which the T_2^{ab} should satisfy

$$\begin{aligned} \delta_{bc}T_2^{ad} - \delta_{ad}T_2^{cb} &= \sum_{i \neq j} I_i^{ab} I_j^{cd} \left\{ \lambda \sum_{k,l} I_i^{kl} I_j^{lk} (D_j - D_i) + [D_i, D_j] \right\} \\ &\quad + \sum_i (\delta_{bc} I_i^{ad} D_i^2 - \delta_{ad} I_i^{cb} D_i^2) \ . \end{aligned} \tag{4.45}$$

A sufficient solution of eq. (4.45) is

$$T_2^{ab} = \sum_i I_i^{ab} D_i^2 \tag{4.46}$$

with

$$[D_i, D_j] = \lambda \sum_{a,b} I_j^{ab} I_i^{ba} (D_i - D_j) \ . \tag{4.47}$$

Thus eq. (4.41) generates long-range interaction through the eq. (4.44) and (4.47). However so far there is not simple relationship between D_i and I_j^{ab} which should satisfy eq. (4.44). It is very difficult to determine the general relationship. Fortunately, BGHP [17] have set up the link with the help of projection. Let the permutation groups Σ_1, Σ_2 and Σ_3 be generated by K_{ij}, P_{ij} and the product $P_{ij}K_{ij}$ respectively, where K_{ij} exchange the positions of particles and P_{ij} exchange the spins at position i and j. The projection ρ was defined as

$$\rho(ab) = a \quad \text{for } \forall a \in \Sigma_2, b \in \Sigma_1 \ , \tag{4.48}$$

i.e. the wave function considered is symmetric. Let I_i^{ab} be the fundamental representations, then

$$P_{ij} = \sum_{a,b} I_i^{ab} I_j^{ba} \ . \tag{4.49}$$

Suppose that there exists [17]

$$D_i = \rho(\hat{D}_i), \quad D_i \in \Sigma_2, \ \hat{D}_i \in \Sigma_1 \tag{4.50}$$

and the $\hat{D}_i$ is particle-like operators, i.e.

$$K_{ij}\hat{D}_i = \hat{D}_j K_{ij}, \quad K_{ij}\hat{D}_l = \hat{D}_l K_{ij} \quad (l \neq i, j) \ . \tag{4.51}$$

Define

$$T_m^{ab} = \sum_i I_i^{ab} \rho(\hat{D}_i^m) \quad (m \geq 0) \ , \tag{4.52}$$

then

$$\text{(a)} \qquad [\hat{D}_j, \hat{D}_i] = \lambda \rho^{-1}(P_{ij}(D_j - D_i)) = \lambda(\hat{D}_j - \hat{D}_i)K_{ij} \ . \tag{4.53}$$

(b) T_m^{ab} satisfy eq. (4.36), i.e., RTT relation eq. (4.32). Actually $f_1^n = 0$ is easy to be checked. By using

$$[\hat{D}_i^n, \hat{D}_j^m] = \sum_{k=0}^{n-1} \hat{D}_i^k [\hat{D}_i, \hat{D}_j^m] \hat{D}_j^{n-k-1} = \lambda \sum_{k=0}^{n-1} \hat{D}_i^k (\hat{D}_i^m - \hat{D}_j^m) \hat{D}_j^{n-k-1} K_{ij} \ ,$$

we have $f_3^{n,m} = 0$.

The projection procedure is very important for it enables us to prove that eq. (4.36) is satisfied by virtue of eq. (4.50).

With the expansion eqs. (4.33) and the projected long-range expansion eq. (4.52), the hamiltonian associated to $T(u)$ is obtained by the expansion of the deformed determinant [17]:

$$\mathrm{det}_q T(u) = \sum_\sigma \epsilon(\sigma) T_{1\sigma_1}(u-(p-1)\lambda) T_{2\sigma_2}(u-(p-2)\lambda)\cdots T_{p\sigma_p}(u)\ . \tag{4.54}$$

A calculation gives

$$\begin{aligned}\mathrm{det}_q T(u) \;=\;& 1+\frac{\lambda}{u}M+\frac{\lambda}{u^2}\left[\rho(\sum_i \hat{D}_i-\frac{\lambda}{2}\sum_{j\neq i}K_{ij})+\frac{\lambda}{2}M(M-1)\right] \\ &+\frac{\lambda}{u^3}\rho\left\{(\sum_i \hat{D}_i-\frac{\lambda}{2}\sum_{j\neq i}K_{ij})^2+\frac{\lambda^2}{12}\sum_{i\neq j\neq k\neq i}K_{ij}K_{jk}\right. \\ &+\lambda(M-1)\sum_i(\hat{D}_i-\frac{\lambda}{2}\sum_{j\neq i}K_{ij}) \\ &\left.+\frac{\lambda^2}{6}M(M-1)(M-2)+\frac{\lambda^2}{4}M(M-1)\right\}+\cdots\ .\end{aligned} \tag{4.55}$$

One takes the Hamiltonian as

$$H=\frac{1}{2}\rho\left\{(\sum_i \hat{D}_i-\frac{\lambda}{2}\sum_{i\neq j}K_{ij})^2+\frac{\lambda^2}{12}\sum_{i\neq j\neq k\neq i}K_{ij}K_{jk}\right\}\ . \tag{4.56}$$

Therefore we define the Hamiltonian which have the *Yangian* symmetry given by eqs.(4.52), (4.42) and (4.47). In comparison to the known models we list the expressions for $\hat{D}_i$ satisfying eq. (4.53) [18]

(1) $\hat{D}_i = p_i+\frac{\lambda}{2}\sum_{i\neq j}[\mathrm{sgn}(x_i-x_j)+1]K_{ij},\quad \lambda=2il,$

$$H=\frac{1}{2}\sum_i p_i^2+\frac{1}{2}\sum_{i\neq j} l(l-P_{ij})\delta(x_i-x_j)\ . \tag{4.57}$$

(2) $\hat{D}_i = p_i+\sum_{i\neq j} l[i\cot a(x_i-x_j)+1]K_{ij},\ \lambda=2l,$

$$H=\frac{1}{2}\sum_i p_i^2+\frac{1}{2}\sum_{i\neq j}\frac{l(l-aP_{ij})}{\sin^2 a(x_i-x_j)}\ . \tag{4.58}$$

(3) $\hat{D}_i = p_i+il\sum_{i\neq j}[\coth a(x_i-x_j)P_{ij}^+ + \tanh a(x_i-x_j)P_{ij}^- +1]K_{ij},\ \lambda=2il\ ,$

$$H=\frac{1}{2}\sum_i p_i^2+\frac{1}{2}\sum_{i\neq j} l(l-aP_{ij})\left(\frac{P_{ij}^+}{\sinh^2 a(x_i-x_j)}-\frac{P_{ij}^-}{\cosh^2 a(x_i-x_j)}\right). \tag{4.59}$$

Eqs. (4.57) and (4.58) were given in Ref. [18], eq. (4.58) was studied in Ref. [17]. Eq (4.59) is the generalization of S-R model.

An alternative description of transfer matrix was given by BGHP [17]. Define

$$\bar{D}_i = \hat{D}_i - \lambda \sum_{i<j} K_{ij} \ , \tag{4.60}$$

then

$$[\bar{D}_i, \bar{D}_j] = 0 \ , \tag{4.61}$$

$$[K_{ij}, \bar{D}_k] = 0 \ \ (k \neq i, j) \ , \tag{4.62}$$

$$K_{ij}\bar{D}_i - \bar{D}_j K_{ij} = \lambda \ . \tag{4.63}$$

It was proved that

$$\bar{T}_i(u) = 1 + \lambda \frac{P_{0i}}{u - \bar{D}_i}, \ \ \bar{T}(u) = \prod_i \bar{T}_i(u) \text{ and } \rho(\bar{T}(u)) \tag{4.64}$$

all satisfy the RTT relation.

The deformed determinant of $\bar{T}(u)$ was defined by

$$\det{}_q \bar{T}(u) = \frac{\Delta_M(u+\lambda)}{\Delta_m(u)} \ , \ \ \Delta_M(u) = \prod_{i=1}^{M} (u - \bar{D}_i) \ . \tag{4.65}$$

It was proved that

$$\rho(\det{}_q \bar{T}(u)) = \det{}_q(T(u)) \ . \tag{4.66}$$

To contain the model eq.(4.28), we define $\bar{D}_i$ related to the $\bar{\pi}_i$ given by eq. (4.21) as

$$\bar{D}_i = \bar{\pi}_i - \beta \sum_{j<i} P^{+}_{ij} K_{ij} \tag{4.67}$$

which satisfies eqs.(4.61), (4.62) and (4.64) etc. So we can put the models eqs. (4.12) and (4.28) into Yang-Baxter system.

In conclusion of this section we have shown the consistence between *Yangian* symmetry and the integrability of Polychronakos for long-range interaction models [18] and given the interpretation of S-R model [19, 20] and Yan model [21] from the point of view of YB system.

§5 Truncated Transfer Matrix $T(u)$ and G-C Top

For the R-matrix is given by eq. (1.15) the case (A) in the section II yields *Yangian*. Now let's look for what kind of algebra appears for the case (B), i.e., $T^{(0)} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. In this section the transfer matrix for the case (B) is denoted by $t(u)$, the RTT relation is calculated to give the following relations($a, b = 1, 2$).

$$[t^{(m)}_{ab}, t^{(k)}_{ab}] = 0, \ \ [t^{(m)}_{ab}, t^{(k)}_{cd}] = [t^{(k)}_{ab}, t^{(m)}_{cd}], \ (m, k \geq 1) \tag{5.1}$$

that are the same as those for the case (A), the other relations are given by

$$\left\{\begin{array}{l} [t_{11}^{(1)}, t_{22}^{(k)}] = [t_{12}^{(1)}, t_{22}^{(k)}] = [t_{21}^{(1)}, t_{22}^{(k)}] = 0, \\ t_{12}^{(k)} = [t_{12}^{(k)}, t_{11}^{(1)}],\ t_{21}^{(k)} = -[t_{21}^{(k)}, t_{11}^{(1)}],\ t_{22}^{(k)} = [t_{12}^{(k)}, t_{21}^{(1)}],\ (k \geq 1) \end{array}\right. \tag{5.2}$$

$$\left\{\begin{array}{l} [t_{22}^{(2)}, t_{11}^{(k)}] + t_{21}^{(k)} t_{12}^{(1)} - t_{21}^{(1)} t_{12}^{(k)} = 0, \\ [t_{12}^{(2)}, t_{22}^{(k)}] + t_{12}^{(k)} t_{22}^{(1)} - t_{12}^{(1)} t_{22}^{(k)} = 0,\ (k \geq 1) \\ [t_{21}^{(2)}, t_{22}^{(k)}] + t_{22}^{(k)} t_{21}^{(1)} - t_{22}^{(1)} t_{21}^{(k)} = 0 \end{array}\right. \tag{5.3}$$

and the iteration relations

$$\begin{array}{rcl} t_{12}^{(k+1)} & = & [t_{12}^{(2)}, t_{11}^{(k)}] + t_{11}^{(k)} t_{12}^{(1)} - t_{11}^{(1)} t_{12}^{(k)}, \\ t_{21}^{(k+1)} & = & [t_{11}^{(k)}, t_{21}^{(2)}] + t_{11}^{(k)} t_{21}^{(1)} - t_{11}^{(1)} t_{21}^{(k)},\ (k \geq 2) \\ t_{22}^{(k+1)} & = & [t_{12}^{(2)}, t_{21}^{(k)}] + t_{11}^{(k)} t_{21}^{(1)} - t_{11}^{(1)} t_{22}^{(k)}. \end{array} \tag{5.4}$$

Not all the above relations are independent. By substituting eq. (5.4) into eq. (5.2) and taking eq. (5.1) into account it can be shown that only $k = 1, 2$ in eq. (5.2) is independent. Thus eq. (5.2) is simplified to

$$\begin{array}{l} [t_{12}^{(k)}, t_{11}^{(1)}] = t_{12}^{(k)},\quad [t_{12}^{(1)}, t_{22}^{(k)}] = 0, \\ [t_{11}^{(1)}, t_{21}^{(k)}] = t_{21}^{(k)},\quad [t_{21}^{(1)}, t_{22}^{(k)}] = 0,\ (k = 1, 2) \\ [t_{12}^{(1)}, t_{21}^{(k)}] = t_{22}^{(k)},\quad t_{22}^{(1)} \text{ is a center.} \end{array} \tag{5.2'}$$

Because $t_{22}^{(1)}$ is a center and

$$C_2 = t_{22}^{(2)} + t_{22}^{(1)} + t_{11}^{(1)} t_{22}^{(1)} - t_{12}^{(1)} t_{21}^{(1)} \tag{5.5}$$

is then also a center:

$$[C_2, t_{ab}^{(k)}] = 0. \tag{5.6}$$

One is able to show that the relation given by eq. (5.3) can be reduced to

$$[t_{22}^{(2)}, t_{11}^{(k)}] + t_{21}^{(k)} t_{12}^{(1)} - t_{21}^{(1)} t_{12}^{(k)} = 0,\ (k \geq 2). \tag{5.3'}$$

Therefore now eqs. (5.1), (5.2'), (5.3') and (5.4) are independent relations. The simplest realization of eq. (5.2') for $k = 1$ is

$$t^{(1)} = \left[\begin{array}{cc} M & \bar{P} \\ P & C \end{array}\right] \tag{5.7}$$

where C is a center and

$$[M, P] = P,\quad [M, \bar{P}] = -\bar{P},\quad [P, \bar{P}] = C \tag{5.8}$$

A simple realization of eq. (5.7) is when $C = 0$:

$$t^{(1)} = \left[\begin{array}{cc} ip & e^{-q} \\ e^{q} & 0 \end{array}\right] \tag{5.9}$$

where

$$[p, q] = -i \tag{5.10}$$

i.e., p and q obey the Heisenberg algebra.

To determine $t^{(2)}$ we should find a realization satisfying eqs. (5.1) , (5.2') and (5.3') for $k = 2$. The corresponding relations for $k > 2$ yield the constraints to the considered operators. This will gives rise to a new type of infinite algebra. However what we are interested in is to find a more physical example where the expansion of $t(u)$ is truncated

$$t(u) = \sum_{n=0}^{m} u^{-n} t^{(n)} \tag{5.11}$$

i.e.,

$$t^{(m+1)} = 0, \quad (\text{all } t^{(k)} = 0 \text{ for } k \geq m+1). \tag{5.12}$$

This possibility is prohibited for the case (A) which leads to *Yangian*, however, eq. (5.12) can be allowed by $t^{(0)} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. This fact can be verified by direct calculation. Let us look for an exact solution of $m = 3$, i.e., by substituting

$$t(u) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + u^{-1} t^{(1)} + u^{-2} t^{(2)} + u^{-3} t^{(3)} \tag{5.13}$$

into the RTT relation to find a sufficient solution of $t^{(1)}$, $t^{(2)}$ and $t^{(3)}$. This solution is related to the Goryachev-Chaplegin gyrostat discussed in Refs. [22, 23].

This model is interesting for the finite terms in the expansion of $t(u)$. The truncation of $t(u)$ corresponds to the finite number of conserved quantities and the satisfaction of RTT relation indicates the quantum integrability of the model.

The Goryachev-Chaplygin(G-C) top is completely integrable in sense of Liouville as shown in the literature [22]. It describe an axially symmetric top moving in an applied field. In the quantum case the G-C top is extended to the quantum G-C gyrostat whose Hamiltonian takes the form:

$$H = \frac{1}{2}(J^2 + 3J_3^2) - bx_1 + pJ_3 \tag{5.14}$$

where $J^2 = J_1^2 + J_2^2 + J_3^2$ and b a parameter. We should note that in this section the notations J_i express the angular momentum. They are irrelevant to the J_λ-operators for *Yangian* shown before. The quantities appearing in eq. (5.14) satisfy the following commutation relations:

$$\begin{aligned} [J_i, J_j] &= -i\epsilon_{ijk} J_k \ , \\ [J_i, x_j] &= -i\epsilon_{ijk} x_k \ , \\ [x_i, x_j] &= 0 \ , \\ [p, q] &= -i \ , \\ [p, J_i] &= [p, x_i] = [q, x_i] = [q, J_i] = 0 \end{aligned} \tag{5.15}$$

where $i,\ j,\ k$ take over 1, 2 and 3. The momentum p and coordinate q of mass center commute with all the locally dynamic variables. The Hamiltonian eq. (5.14) commutes with the other integral of motion:

$$G = (2J_3 + p)(J^2 - J_3^2) + 2b[x_3, J_1]_+ \ . \tag{5.16}$$

where $[A, B]_+ \equiv AB + BA$. In addition to the commutation relations eq. (5.15) there are two constraints:

$$\begin{aligned} \sum_{i=1}^{3} J_i x_i &= 0, && (5.17) \\ \sum_{i}^{3} x_i^2 &= 1. && (5.18) \end{aligned}$$

Eq. (5.17) plays important roles in proving the quantum integrability of the model.

Now let us turn to the RTT relation.

For solving eq. (1.7) with the form eq. (5.13) one first put

$$t_{11}^{(1)} = \alpha p, \quad t_{22}^{(1)} = 0, \quad t_{12}^{(1)} = \beta e^{\tau q} x_+, \quad t_{21}^{(1)} = \gamma e^{-\tau q}, \tag{5.19}$$

and

$$\begin{aligned} t_{11}^{(2)} &= f_1 J^2 + f_2 J_3^2 + f_3 p J_3 + f_4 x_- + f_5, \\ t_{11}^{(3)} &= (p + g_3 J_3)(g_1 J^2 + g_2 J_3^2 + g_5) + g_4 [J_-, x_3]_+ && (5.20) \end{aligned}$$

where $f_1, \ldots, f_5,\ g_1, \ldots, g_4,\ \alpha,\ \beta,\ \gamma$ and τ all parameters to be determined and $J_\pm = J_1 \pm iJ_2$, $x_\pm = x_1 \pm ix_2$ obey the commutation relations shown by eq. (5.15). Obviously not all the parameters are independent. λ can be normalized to be one. Substituting eqs. (5.19) and (5.20) together with eq. (5.17) into eq. (1.7) after lengthy calculations by hand we find

$$\begin{aligned} &\tau = -i\alpha^{-1}\lambda,\ f_1 = -\frac{1}{4}\lambda,\ f_2 = -\frac{3}{4}\lambda,\ f_3 = \alpha,\ f_5 = -\frac{1}{16}\lambda, \\ &g_1 = -g_2 = -\frac{1}{4}\alpha,\ g_3 = -\lambda\alpha^{-1},\ g_4 = \frac{1}{4}f_4,\ g_5 = -\frac{1}{16}\alpha. \end{aligned}$$

Denoting $f_4 = f$ the solution of $t^{(n)}$ reads:

$$\begin{aligned} t_{11}^{(2)} &= -\frac{1}{4}\lambda(J^2 + 3J_3^2 + \frac{1}{4}) + \alpha p J_3 + f x_-, \quad t_{22}^{(2)} = \lambda^{-1}\gamma\beta x_+, \\ t_{12}^{(2)} &= -\lambda^{-1}\beta e^{\tau q}\left(-\frac{\lambda}{4}[J_+, x_3]_+ + x_+(\lambda J_3 - \alpha p)\right), \quad t_{21}^{(2)} = \gamma e^{-\tau q} J_3, && (5.21) \\ t_{11}^{(3)} &= -\frac{1}{4}\alpha(p - \lambda\alpha^{-1}J_3)(J^2 - J_3^2 + \frac{1}{4}) + \frac{1}{4}f[J_-, x_3]_+, \\ t_{22}^{(3)} &= \frac{1}{4}\lambda^{-1}\beta\gamma[J_+, x_3]_+, \\ t_{12}^{(3)} &= -\lambda^2\beta e^{\tau q}\left\{f x_3^2 - \frac{1}{4}\alpha[J_+, x_3]_+(p - \lambda\alpha^{-1}J_3)\right\}, \\ t_{21}^{(3)} &= -\frac{1}{4}\gamma e^{-\tau q}(J^2 - J_3^2 + \frac{1}{4}). && (5.22) \end{aligned}$$

Therefore the truncated *Yangian* can be viewed as a mapping of the algebra given by eq. (5.15).

Substituting the derived $t(u)$ into eq. (2.58) it follows

$$\mathrm{det}t(u) = u^{-3}(u-1)^{-3}\lambda^{-1}f(u^2 - u + \frac{3}{16})(x_+x_- + x_3^2). \tag{5.23}$$

So only $\sum_{i=1}^3 x_i^2$ is a Casimir operator, i.e. $C_m = 0$ for $m < 4$, $C_4 = \lambda^{-1}f\sum_{i=1}^3 x_i^2$ that commute with J_i. The constraint eq. (5.18) does not play role in solving the RTT relation. By choosing a proper representation eq. (5.18) can be taken that is automatically satisfied by virtue of eq. (2.52). Notice that $\mathrm{det}t(u)$ has zero's at $u = \frac{1}{4}$ and $u = \frac{3}{4}$ where the inverse of $t(u)$ can not be defined.

It is worth noting that the commutation relations shown by eq. (5.15) are invariant subject to a transformation:

$$J'_a = \sum_b A_{ab}J_b,\ x'_a = \sum_b A_{ab}x_b, \quad (a, b = 1, 2) \tag{5.24}$$

where

$$\begin{array}{ll} A_{11} = \epsilon A_{22}, & A_{12} = -\epsilon A_{21}, \\ A_{11}^2 + A_{21}^2 = 1, & \epsilon = \pm 1 = \mathrm{det}A \end{array}. \tag{5.25}$$

This transformation is useful to transform the Hamiltonian in preserving the RTT relation.

Let us turn to the conserved quantities for G-C gyrostat.

By taking the trace of the transfer matrix one obtains

$$\begin{aligned} \mathrm{tr}t(u) \quad = \quad & \lambda + u^{-1}\alpha p + u^{-2}\left\{-\frac{1}{4}\lambda(J^2 + 3J_3^2 + \frac{1}{4}) + \alpha pJ_3 + fx_- + \lambda^{-1}\beta\gamma x_+\right\} + \\ & \frac{u^{-3}}{4}\left\{-(\alpha p - \lambda J_3)(J^2 - J_3^2 + \frac{1}{4}) + f[J_-, x_3]_+ + \lambda^{-1}\beta\gamma[J_+, x_3]_+\right\}. \end{aligned} \tag{5.26}$$

Hence we have the Hamiltonian H_p and the other conserved quantities G_p:

$$H_p = \frac{1}{2}\left\{\frac{1}{4}\lambda(J^2 + 3J_3^2) - fx_- - \lambda^{-1}\beta\gamma x_+ - \alpha pJ_3\right\} + \frac{1}{16}\lambda, \tag{5.27}$$

$$G_p = \frac{1}{4}\left\{(\alpha p - \lambda J_3)(J^2 - J_3^2 + \frac{1}{4}) + f[J_-, x_3]_+ + \lambda^{-1}\beta\gamma[J_+, x_3]_+\right\}. \tag{5.28}$$

If one requires

$$4\lambda^{-1}\beta\gamma f = b^2, \tag{5.29}$$

then there is rotational invariance about the x_3 axis. By virtue of eqs. (5.22) and (5.24) we find that eqs. (5.27) and (5.28) are reduced to

$$H_p \quad = \quad \frac{1}{2}\left\{\frac{1}{4}\lambda(J'^2 + 3J'^2_3) - bx'_1 - \alpha p\epsilon J'_3\right\} + \frac{1}{16}\lambda, \tag{5.30}$$

$$G_p \quad = \quad -\frac{1}{4}\left\{(\alpha p - \epsilon\lambda J'_3)(J'^2 - J'^2_3 + \frac{1}{4}) + b[x_3, J'_1]_+\right\} \tag{5.31}$$

which are exactly those given by Sklyanin [23]. Obviously, the parameters λ and τ are trivial in the solution of $t(u)$ and β, γ can be viewed as the consequence of $t(u)$ subject to a similar transformation. Therefore only parameters f and α are essential in determining the form of $t(u)$ though f does not appear in the Hamiltonian. The simplest realization of quantum G-C top is a quantum mechanical system on sphere.

Taking the constraints eqs. (5.17) and (5.18) into account the simplest realization of J_i and x_i can be made through

$$\begin{aligned} x_i &= n_i(t), && (5.32)\\ J_i &= -i\epsilon_{ijk}n_j(t)\dot{n}_k(t)\ (i,j,k=1,2,3) && (5.33) \end{aligned}$$

that give $J^2 = -\dot{\vec{n}}\cdot\dot{\vec{n}}$.

In conclusion we have shown that the truncated *Yangian* gives rise to quantum integrable system with finite number of conserved quantities and make the system work in the space higher than (1+1) dimensions.

The trigonometric extension of the truncated *Yangian* will be the "truncated" affine quantum algebra which is the generalization of Drinfeld's statement discussed in Ref. [25, 26]. It gives the q-deformed G-C gyrostat which will be discussed in the next section.

§6 Trigonometric G-C Gyrostat

If R-matrix is taken to be the simple 6-vertex from (1.17), the trigonometric transfer matrix $T(x)$ can be found on the basis of RTT relation, where $x = e^u$(or $x = e^{iu}$) is the spectral parameter. In terms of x, the RTT relation takes the form

$$\check{R}(xy^{-1})(T(x)\otimes T(y)) = (T(y)\otimes T(x))\check{R}(xy^{-1}) \tag{6.1}$$

$T(x)$ can be expanded as

$$T(x) = \sum_{n=-\infty}^{+\infty} x^n T^{(n)}. \tag{6.2}$$

A simple form of trigonometric R-matrix(which is a deformation of eq. (1.18)) is given by

$$\check{R}(u) = \begin{bmatrix} \sin(u+\eta) & & & \\ & e^{-i\theta}\sin\eta & \sin u & \\ & \sin u & e^{i\theta}\sin\eta & \\ & & & \sin(u+\eta) \end{bmatrix} \tag{6.3}$$

where $x = e^{iu}$, η and θ are constants. Set $q = e^{-i\eta}$, eq. (6.3) can be recast into

$$\check{R}(x) = x\check{R} - x^{-1}\check{R}^{-1} \tag{6.4}$$

where

$$\check{R} \;=\; \begin{bmatrix} q^{-1} & & & \\ & 0 & 1 & \\ & 1 & q^{-1}-q & \\ & & & q^{-1} \end{bmatrix} \tag{6.5}$$

which is the asymptotic form of $\check{R}(x)$ as $x \to \infty$. It can be verified that there is a subset in $T(x)$:

$$T(x) = xT_{+} - x^{-1}T_{-} \tag{6.6}$$

satisfying eq. (6.2) for the $\check{R}$ given by eq. (6.3). Actually, substituting eq. (6.5) into eq. (6.3) it follows [24]:

$$\begin{array}{l} \check{R}(T_{\pm} \otimes T_{\pm}) = (T_{\pm} \otimes T_{\pm})\check{R} \\ \check{R}(T_{+} \otimes T_{-}) = (T_{-} \otimes T_{+})\check{R} \end{array} \tag{6.7}$$

where $\check{R}$ is given by eq. (6.5). It is noted that each element of the matrices $T_{\pm}$ is quantum operator. One can check that

$$T_{+} = \begin{bmatrix} K & (q^{-1}-q)X^{+} \\ 0 & K^{-1} \end{bmatrix}, \quad T_{-} = \begin{bmatrix} K^{-1} & 0 \\ (q^{-1}-q)X^{-} & K \end{bmatrix} \tag{6.8}$$

satisfy eq. (6.7) when $X^{\pm}$ and K satisfy

$$KX^{\pm}K^{-1} = q^{\pm}X^{\pm}, \quad [X^{+}, X^{-}] = \frac{K^{2} - K^{-2}}{q - q^{-1}} \tag{6.9}$$

that is known as the quantum algebra associated with $SU(2)$, denoted by $U_q(SU(2))$. The pioneered works concerning quantum algebras were made by Drinfeld [25, 26], Jimbo [27] and Faddeev [28] although the simplest form eq. (6.9) was first presented by Kulish and Sklyanin [5].

The general form of trigonometric $T(x)$ is very complicated and denoted by $U_q(\widehat{SU(2)})$, i.e., the q-affine algebra [25]. The comparison between $Y(sl(2))$ and $U_q(sl(2))$ is shown by the following chart-follow ($w = q - q^{-1}$):

$$\begin{tikzcd}
\check{R} = uP + I \arrow[dr] & & \check{R}\text{ given by eq. (6.3)} \arrow[ll, "q=e^{-i\eta}", "\begin{array}{c}\text{rational}\\ \text{limit}\end{array}"'] \arrow[dl] \\
& \boxed{\text{RTT relation}} \arrow[dl] \arrow[dr] & \\
T(u) = \sum_{n=0}^{\infty} u^{-n}T^{(n)} & & \begin{array}{c} T(x) = \sum_{n=-\infty}^{+\infty} x^{n}L^{(n)} \\ = L^{(0)} + xL^{(1)} + x^{-1}L^{(-1)} + \cdots \\ L^{(1)} = L_{+},\ L^{(-1)} = -L_{-} \end{array}
\end{tikzcd}$$

$$\begin{array}{ccc}
T^{(1)} = \left[\begin{array}{cc} I_3 & I_+ \\ I_- & -I_3 \end{array}\right] & \begin{array}{c}\text{rational} \\ \longleftarrow \\ \text{limit}\end{array} & \begin{array}{c} L_+ = \left[\begin{array}{cc} K & wX^+ \\ 0 & K^- \end{array}\right], \\ L_- = \left[\begin{array}{cc} K^{-1} & 0 \\ -wX^- & K \end{array}\right] \end{array} \\
J_\lambda,\ T^{(n)}(n \geq 3) & & \sum_n x^n L^{(n)} \\
Y(sl(2)) & & U_q(sl(2)) \\
\text{deformed loop algebra} & & q\text{-deformed Kac-Moody algebra}
\end{array}$$

To find such $T(x)$-matrix let us consider 2 by 2 matrix, substitute

$$T(x) = \left[\begin{array}{cc} T_{11}(x) & T_{12}(x) \\ T_{21}(x) & T_{22}(x) \end{array}\right] = \sum_{n=-\infty}^{+\infty} x^n T^{(n)} \tag{6.10}$$

or

$$T_{ab}(x) = \sum_{n=-\infty}^{+\infty} x^n T_{ab}^{(n)}\ (a, b = 1, 2) \tag{6.11}$$

into eq. (6.1) we derive

$$\begin{array}{l}
[T_{ab}^{(k)}, T_{ab}^{(j)}] = 0, \quad [T_{11}^{(k)}, T_{22}^{(j)}] = [T_{11}^{(j)}, T_{22}^{(k)}], \quad [T_{12}^{(k)}, T_{21}^{(j)}] = [T_{12}^{(j)}, T_{21}^{(k)}], \\
[T_{22}^{(k-1)}, T_{11}^{(j+1)}] - [T_{22}^{(k+1)}, T_{11}^{(j-1)}] = w(T_{12}^{(j)} T_{21}^{(k)} - T_{12}^{(k)} T_{21}^{(j)}), \\
[T_{21}^{(k-1)}, T_{12}^{(j+1)}] - [T_{21}^{(k+1)}, T_{12}^{(j-1)}] = w(T_{11}^{(j)} T_{22}^{(k)} - T_{11}^{(k)} T_{22}^{(j)}, \\
qT_{aa}^{(k-1)} T_{ab}^{(j+1)} - q^{-1} T_{aa}^{(k+1)} T_{ab}^{(j-1)} = T_{ab}^{(j+1)} T_{aa}^{(k-1)} - T_{ab}^{(j-1)} T_{aa}^{(k+1)} + w T_{aa}^{(j)} T_{ab}^{(k)}, \\
qT_{ab}^{(k-1)} T_{aa}^{(j+1)} - q^{-1} T_{ab}^{(k+1)} T_{aa}^{(j-1)} = T_{aa}^{(j+1)} T_{ab}^{(k-1)} - T_{aa}^{(j-1)} T_{ab}^{(k+1)} + w T_{ab}^{(j)} T_{aa}^{(k)}, \\
qT_{ba}^{(j+1)} T_{aa}^{(k-1)} - q^{-1} T_{ba}^{(j-1)} T_{aa}^{(k+1)} = T_{aa}^{(k-1)} T_{ba}^{(j+1)} - T_{aa}^{(k+1)} T_{ba}^{(j-1)} + w T_{ba}^{(k)} T_{aa}^{(j)}, \\
qT_{aa}^{(j+1)} T_{ba}^{(k-1)} - q^{-1} T_{aa}^{(j-1)} T_{ba}^{(k+1)} = T_{ba}^{(k-1)} T_{aa}^{(j+1)} - T_{ba}^{(k+1)} T_{aa}^{(j-1)} + w T_{aa}^{(j)} T_{ba}^{(k)}.
\end{array} \tag{6.12}$$

The inverse of $T(x)$ is given by

$$[T(x)]^{-1} = [\det_q T(x)]^{-1} \left[\begin{array}{cc} T_{22}(q^{-1}x) & -T_{12}(q^{-1}x) \\ -T_{21}(q^{-1}x) & T_{11}(q^{-1}x) \end{array}\right] \tag{6.13}$$

where

$$\det_q T(x) = T_{11}(x) T_{22}(q^{-1}x) - T_{12}(x) T_{21}(q^{-1}x) \tag{6.14}$$

It is easy to know that

$$[\det_q T(x), T_{ab}(y)] = 0 \tag{6.15}$$

Correspondingly

$$\det_q T(x) = \sum_{n=-\infty}^{+\infty} x^{-n} C_n, \quad C_n = \sum_{k+j=n} q^{-j} (T_{11}^{(k)} T_{22}^{(j)} - T_{12}^{(k)} T_{21}^{(j)}) \tag{6.16}$$

and

$$\mathrm{tr}T(x)=\sum_{n=-\infty}^{+\infty}(T_{11}^{(n)}+T_{22}^{(n)})x^n \tag{6.17}$$

Definition $T^{(n)}$ is called truncated q-affine quantum algebra if $T^{(n)}=0$ for $|n|\geq 4$.

To solve (6.12) we take the ansatz (motivated by the rational correspondence)

$$\begin{aligned} T_{11}^{(\pm 2)} &= T_{11}^{(0)}=T_{22}^{(\pm 3)}=T_{22}^{(\pm 2)}=T_{22}^{(0)} \\ &= T_{12}^{(\pm 3)}=T_{12}^{(\pm 1)}=T_{21}^{(\pm 3)}=T_{21}^{(\pm 1)}=0 \end{aligned} \tag{6.18}$$

and set

$$T_{11}^{(\pm 3)}=\pm\lambda_3 e^{\mp i\eta P},\quad T_{11}^{(\pm 1)}=A_{\pm}^{(0)}+e^{\mp i\eta P}A_{\pm}^{(1)}+e^{\pm i\eta P}A_{\pm}^{(2)} \tag{6.19}$$

$$T_{12}^{(\pm 2)}=e^{i\xi Q\mp i\eta P}E_{\pm 2},\quad T_{12}^{(0)}=e^{i\xi Q}(E^{(0)}+e^{-i\eta P}E^{(-)}+e^{i\eta P}E^{(+)}) \tag{6.20}$$

$$T_{21}^{(\pm 2)}=\alpha_2 K^{\pm 1}e^{-i\xi Q},\quad T_{21}^{(0)}=e^{-i\xi Q}F_0,\quad q=e^{i\xi\eta} \tag{6.21}$$

where $\lambda_3,\ \alpha_2,\ \xi$ and η are constants, P and Q satisfy

$$[P,Q]=-i \tag{6.22}$$

here $A_{\pm}^{(i)}(i=0,1,2),\ E_{\pm 2},\ E^{(j)}(j=0,\pm),\ F_0$ and K are operators commuting with P and Q. They will be determined by (6.12).

After tedious calculation we find that the following algebraic relations solve the RTT relation for 6-vertex form of R-matrix:

$$\begin{aligned} &KE_{\pm 2}=q^{-1}E_{\pm 2}K \\ &A_{\pm}^{(2)}=\mp\lambda_3 K^{\pm 2},\quad A_{\pm}^{(1)}=\pm\lambda_3\alpha_2^{-1}K^{\mp}S,\quad F_0=S \\ &E^{\pm}=-E_{\pm 2}K^{\pm 2},\quad T_{22}^{(\pm 1)}=\pm\lambda_3^{-1}\alpha_2 E_{\pm 2}K^{\pm 1} \end{aligned} \tag{6.23}$$

where $A_{\pm}^{(0)},\ E_{\pm 2},\ E^{(0)}$ and S satisfy

$$\begin{aligned} &q^2E_{-2}E_{+2}=E_{+2}E_{-2},\quad [A_{+}^{(0)},A_{-}^{(0)}]=[K,S]=0 \\ &KA_{\pm}^{(0)}K^{-1}=qA_{\pm}^{(0)},\quad q^{\pm 1}E_{\pm 2}A_{\mp}^{(0)}=A_{\mp}^{(0)}E_{\pm 2} \\ &q^{\pm 1}E_{\pm 2}A_{\pm}^{(0)}-A_{\pm}^{(0)}E_{\pm 2}=\lambda_3 wE^{(0)} \\ &SA_{\pm}^{(0)}-q^{\mp 1}A_{\pm}^{(0)}S=\pm\alpha_2 wA_{\mp}^{(0)}K^{\pm 1} \\ &SE_{\pm 2}-q^{\pm 1}E_{\pm 2}S=\pm\alpha_2 wE_{\mp 2}K^{\mp 1} \end{aligned} \tag{6.24}$$

A realization of the algebra eqs. (6.2) and (6.24), i.e., $T^{(n)}(T^{(n)}=0$ for $|n|\geq 4)$, can be made by

$$\begin{aligned} &A_{\pm}^{(0)}=\lambda_{\pm}^{(1)}\hat{J}_-\hat{x}_3+\lambda_{\pm}^{(2)}\hat{x}_-, \\ &E_{\pm 2}=\beta_{\pm}^{(1)}\hat{J}_+\hat{x}_3+\beta_{\pm}^{(2)}\hat{x}_+, \\ &S=\lambda_1\hat{J}_+\hat{J}_-+\lambda_2, \\ &K=\exp(i\xi\eta\hat{J}_3) \end{aligned} \tag{6.25}$$

where $E^{(0)},\ \lambda_{\pm}^{(i)},\ \beta_{\pm}^{(i)}$ and $\lambda_i (i=1,2)$ are K-dependent operators which are given by

$$\alpha_2^2 = g\alpha^2, \quad \lambda_1 = (g-1)wg^{-1}\alpha,\ \lambda_2 = \alpha(q-K+K^{-1}), \tag{6.26}$$

and

$$\begin{aligned}
\lambda_-^{(1)} &= 0,\ \lambda_+^{(1)} = \lambda^{(+)}K^{2(1-\delta_+)-\beta},\ \lambda_+^{(2)} = (1-q^{-1})^{-1}q^{-\delta_+}\tau_-K^{-1+\delta_+}\lambda_+^{(1)} \\
\lambda_-^{(2)} &= (1-q^{-1})^{-1}q^{-1-\delta_+}\alpha^{-1}\alpha_2\tau_-K^{-1+\delta_+}\lambda_+^{(1)}, \quad \beta_+^{(1)} = 0, \\
\beta_-^{(2)} &= -(1-q^{-1})^{-1}q^{-1+\delta_+}\tau_+K^{\delta_+}\beta_-^{(1)}, \quad \beta_-^{(1)} = \beta^{(-)}K^{\beta}, \\
\beta_+^{(2)} &= (1-q^{-1})^{-1}q^{-4+\delta_+}\alpha^{-1}\alpha_2\tau_+K^{\delta_+-2}\beta_-^{(1)}, \\
E^{(0)} &= -\lambda_3^{-1}w^{-1}(1-q^{-1})^{-1}q^{-3+\delta_+}g\alpha^{-1}\alpha_2K^{-1}\beta_-^{(1)}\lambda_+^{(1)}(qK)(\hat{x}_3)^2
\end{aligned} \tag{6.27}$$

for $\delta_+ + \delta_- = 1$.

$$\begin{aligned}
\lambda_+^{(1)} &= 0,\ \ \lambda_{(-)}^{(1)} = \lambda^{-}K^{2(1+\delta_+)-\beta},\ \lambda_-^{(2)} = (1-q^{-1})^{-1}q^{-\delta_+}\tau_-K^{1+\delta_+}\lambda_-^{(1)}, \\
\lambda_+^{(2)} &= (1-q^{-1})^{-1}q^{-\delta_+}\alpha^{-1}\alpha_2\tau_-K^{1+\delta_+}\lambda_-^{(1)}, \quad \beta_-^{(1)} = 0, \\
\beta_+^{(2)} &= -(1-q^{-1})^{-1}q^{1+\delta_+}\tau_+K^{\delta_+}\beta_+^{(1)}, \quad \beta_+^{(1)} = \beta^{(+)}K^{\beta}, \\
\beta_-^{(2)} &= (1-q^{-1})^{-1}q^{4+\delta_+}\alpha\alpha_2^{-1}\tau_+K^{\delta_++2}\beta_+^{(1)}, \\
E^{(0)} &= -\lambda_3^{-1}w^{-1}(1-q^{-1})^{-1}q^{3+\delta_+}g\alpha\alpha_2^{-1}K\beta_+^{(1)}\lambda_-^{(1)}(qK)(\hat{x}_3)^2
\end{aligned} \tag{6.28}$$

for $\delta_+ + \delta_- = -1$, where $\delta_+,\ \alpha,\ \beta^{(\pm)},\ \lambda^{(\pm)}$ and $\tau_\pm$ are arbitrary constants. It is noted that the parameters $\beta_{\mp}^{(1)}(K)$ depend on K and $\lambda_{\pm}^{(1)}(qK)$ on qK, respectively, whereas $\delta_+,\ \alpha,\ \beta^{(\pm)},\ \lambda^{(\pm)}$ and $\tau_\pm$ are arbitrary constants.

The operators $\hat{J}_\pm,\ \hat{J}_3,\ \hat{x}_\pm$ and $\hat{x}_3$ in eq. (6.25) obey the following relations

$$\begin{aligned}
[\hat{J}_\pm, \hat{x}_\pm] &= [\hat{J}_3, \hat{x}_3] = 0, \\
[\hat{J}_\pm, \hat{J}_3] &= \pm\hat{J}_3, \quad [\hat{J}_+, \hat{J}_-] = -g[\hat{J}_3]_q, \\
[\hat{x}_\pm, \hat{J}_3] &= \pm\hat{x}_\pm, \\
q^{\delta_\mp}\hat{J}_\pm\hat{x}_3 &= \hat{x}_3\hat{J}_\pm \pm \tau_\pm K^{\pm\delta_\pm}\hat{x}_\pm, \\
[\hat{x}_+, \hat{x}_-] &= 0, \quad \hat{x}_\pm\hat{x}_3 = q^{\delta_\pm}\hat{x}_3\hat{x}_\pm, \\
q^{-1}\hat{J}_\pm\hat{x}_\mp &= \hat{x}_\mp\hat{J}_\pm \mp \tau_\mp^{-1}gK^{\mp\delta_\pm}\hat{x}_3
\end{aligned} \tag{6.29}$$

together with the constraint

$$g[\hat{J}_3]_q\hat{x}_3 + \tau_-K^{-\delta_-}\hat{J}_+\hat{x}_- + \tau_+K^{\delta_+}\hat{J}_-\hat{x}_+ = 0 \tag{6.30}$$

In conclusion eqs. (6.20)-(6.30) solve RTT relation for R is given by eq. (1.15).

The $\det_q T(x)$ for $T^{(n)}(x) = 0(|n| \geq 4)$ can be expressed by $\hat{J}_\pm,\ \hat{x}_\pm$ and $\hat{x}_3$ through

$$\det{}_q T(x) = \{x^2 + q^2x^{-2} + \alpha^{-1}\alpha_2(1+q)\}C_{+2} \quad (\alpha^{-1}\alpha_2 = \pm q^{\frac{1}{2}}) \tag{6.31}$$

where

$$C_{+2} = F_2\{\tau_-K^{\delta_+}(\hat{J}_+\hat{J}_-\hat{x}_3 - \frac{\tau_+K^{\delta_+}\hat{x}_+\hat{x}_-}{1-q} + w^{-1}g(\hat{x}_3)^2)\},\ (w = q - q^{-1}) \tag{6.32}$$

with

$$F_2 = \lambda_3^{-1}\alpha(1-q)^{-1}q^{1-\beta-\delta_+}\beta^{(+)}\lambda^{(-)}K^{2\delta_+} \tag{6.33}$$

that leads to for $\delta_+ + \delta_- = -1$:

$$C_+ = F_1\{\tau_+ K^{\delta_+-1}(\hat{J}_-\hat{J}_+\hat{x}_3 - \frac{\tau_- K^{\delta_+-1}\hat{x}_+\hat{x}_-}{1-q^{-1}} + wg(\hat{x}_3)^2)\} \tag{6.34}$$

where

$$F_1 = \lambda_3^{-1}\alpha(1-q^{-1})^{-1}q^{-2-\beta-\delta_+}\beta^{(-)}\lambda^{(+)}K^{2(1-\delta_+)} \tag{6.35}$$

The non-vanishing C_n in (6.16) are

$$\begin{aligned} C_{\pm 4} &= q^{\mp 1}(T_{11}^{(\pm 3)}T_{22}^{(\pm 1)} - q^{\mp 1}T_{12}^{(\pm 2)}T_{21}^{(\pm 2)}), \\ C_{\pm 2} &= q^{\pm 1}T_{11}^{(\pm 3)}T_{22}^{(\mp 1)} + q^{\mp 1}T_{11}^{(\pm 1)}T_{22}^{(\pm 1)} - T_{12}^{(\pm 2)}T_{21}^{(0)} - q^{\mp 2}T_{12}^{(0)}T_{21}^{(\pm 2)}, \\ C_0 &= qT_{11}^{(+1)}T_{22}^{(-1)} + q^{-1}T_{11}^{(-1)}T_{22}^{(+1)} - q^2T_{12}^{(2)}T_{21}^{(-2)} - q^{-2}T_{12}^{(-2)}T_{21}^{(2)}T_{12}^{(0)}T_{21}^{(0)}. \end{aligned} \tag{6.36}$$

Substituting (6.20)-(6.24) into $\det_q T(x)$ after calculations we obtain $C_{-2} = q^2 C_{+2},\ C_0 = \alpha^{-1}\alpha_2(1+q)C_{+2},\ C_{\pm 4} = 0$ and $C_{+2} = q^{-1}\alpha_2(\lambda_3^{-1}A_+^{(0)}E_{+2} - q^{-1}E^{(0)})K$ that lead to (6.31) and (6.32).

It is straightforward to obtain

$$\begin{aligned} \mathrm{tr}T(x) &= (A_+^{(0)} + e^{-i\eta P}A_+^{(1)} + e^{i\eta P}A_+^{(2)} + \lambda_3^{-1}\alpha_2 E_{+2}K)x \\ &\quad +(A_-^{(0)} + e^{-i\eta P}A_-^{(2)} + e^{i\eta P}A_-^{(1)} + \lambda_3^{-1}\alpha_2 E_{-2}K^{-1})x^{-1} \\ &\quad +\lambda_3(e^{-i\eta P}x^3 - e^{i\eta P}x^{-3}) \end{aligned} \tag{6.37}$$

where $A_\pm^{(0)},\ E_{\pm 2}$ are given by (6.25), whereas $A_\pm^{(1)}$ and $A_\pm^{(2)}$ are given by (6.23). In terms of

$$T(x) = \sum_{n=-3}^{3} x^n T^{(n)} = T(u)\sum_{n,m}\sin^n u\cos^m u T^{(n,m)} \tag{6.38}$$

where $x = e^{iu}$ one finds

$$\det_q T(u) = \{2q\cos(2u - \xi\eta) + \alpha^{-1}\alpha_2(1+q)\}C_{+2} \tag{6.39}$$

and

$$\begin{aligned} \mathrm{tr}T(u) &= T_{11}^{(3,0)}\sin^3 u + T_{11}^{(2,1)}\sin^2 u\cos u \\ &\quad +(T_{11}^{(1,0)} + T_{22}^{(1,0)})\sin u + (T_{11}^{(0,1)} + T_{22}^{(0,1)})\cos u \end{aligned} \tag{6.40}$$

Following the Inverse Scattering Methods $T_{11}^{(3,0)}$ and $T_{11}^{(2,1)}$ correspond to the momentum conservation. The Hamiltonian is defined by

$$H = f_1\{T_{11}^{(1,0)} + T_{22}^{(1,0)}\} \tag{6.41}$$

and another constant of motion is

$$G = f_2\{T_{11}^{(0,1)} + T_{22}^{(0,1)}\} \tag{6.42}$$

where f_1 and f_2 are arbitrary constants.

The Hamiltonian of the system given by eqs. (6.41) and (6.42), take the form

$$\begin{aligned} H &= f_1 \Big(2i\lambda_3\{\alpha\alpha_2^{-1}(q-1)wg^{-1}\cos[\eta(P+\xi\hat{J}_3)]\hat{J}_+\hat{J}_- \\ &\quad +2\sin(\xi\eta\hat{J}_3)(\sin[\eta(P+\xi\hat{J}_3] - \alpha^{-1}\alpha_2 q^{\frac{1}{2}}\sin[\eta(P+\xi\hat{J}_3+\frac{1}{2})]) \\ &\quad + (\alpha\alpha_2^{-1}(q+1)+2)\cos(\xi P)\} + D_+\Big), \end{aligned} \tag{6.43}$$

$$\begin{aligned} G &= f_2 \Big(-2i\lambda_3 \Big\{\alpha\alpha_2^{-1}(q-1)wg^{-1}\hat{J}_+\hat{J}_- + 2(\alpha\alpha_2^{-1}q^{\frac{1}{2}}\cos[\xi\eta(\hat{J}_3+\frac{1}{2})]) \\ &\quad + \cos(\xi\eta\hat{J}_3)\Big\}\sin[\eta(P+\xi\hat{J}_3] + D_-\Big) \end{aligned} \tag{6.44}$$

where $D_\pm$ are given by

$$\begin{aligned} D_\pm &= \epsilon_\pm \Big\{\lambda_+^{(1)}(\hat{J}_-\hat{x}_3 + (1-q^{-1})^{-1}q^{-\delta_+}\tau_- K^{1+\delta_+}(1 \mp q^{-1}\alpha^{-1}\alpha_2)\hat{x}_-) \\ &\quad +\lambda_3^{-1}\alpha_2\beta_+^{(1)}(\pm\hat{J}_\pm\hat{x}_3 K^{-1} + (1-q^{-1})^{-1}q^{-2+\delta_+}\tau_+ \\ &\quad\ K^{\delta_+-1}(\alpha^{-1}\alpha_2 q^{-1} \mp 1)\hat{x}_+)\ \Big\} \end{aligned} \tag{6.45}$$

for $\delta_+ + \delta_- = 1$, and

$$\begin{aligned} D_\pm &= \epsilon_\pm \Big\{\lambda_3^{-1}\alpha_2\beta_+^{(1)}(\hat{J}_\pm\hat{x}_3 K + (1-q)^{-1}q^{2+\delta_+}\tau_+ K^{\delta_++1}(\alpha\alpha_2^{-1}q \mp 1)\hat{x}_+) \\ &\quad +\lambda_-^{(1)}(\mp\hat{J}_+\hat{x}_3 + (1-q)^{-1}q^{-\delta_+}\tau_- K^{1+\delta_+}(\alpha\alpha_2^{-1}q \mp 1)\hat{x}_-)\ \Big\} \end{aligned} \tag{6.46}$$

for $\delta_+ + \delta_- = -1$, with $\epsilon_+ = i,\ \epsilon_- = 1$.

It is noted that there are two possibilities:

$$\alpha = \pm q^{-\frac{1}{2}}\alpha_2 \tag{6.47}$$

in (6.45) and (6.46).

We would like to remark that for the given standard 6-vertex R-matrix eq. (1.18) we find a set of solution for truncated RTT relation. The solution can be realized through the algebra (6.29) and corresponding conserved quantities (6.43) and (6.44). Eq. (6.29) naturally yields the non-commutative geometry [29]. Since the considered system is axially symmetric so that the coordinates on $(x_1,\ x_2)$ plane are commute with each other, whereas the third coordinates x_3 does not commute with them.

It can be shown that the rational limit of the trigonometric conserved quantities H and G given by eqs. (6.43) and (6.44), respectively, are exactly the same as the correspondence in the G-C gyrostat model shown by eqs. (5.27) and (5.28).

All the calculations in this section are tedious and can be found in Ref. [30].

M.-L. Ge and K. Xue would like to thank Prof. Y.M. Cho for his kind invitation, hospitility and the suggestion for the topic of this papers. The authers are indebted to Profs. M. Wadati, Y.S. Wu and B.L. Hao for their stimulating discussions. This work is in part supported by NSF of China.

References

[1] L.I. Schiff, "*Quantum Mechanics*", McGraw-Hill Kogakusha, Tokyo 1969.

[2] H.B. Thacker, Exact integrability in quantum field theory and statistical systems, *Rev. of Mod. Phys.*, **53**(1981)253.

[3] M. Gaudin, "*La Fonction d'onde de Bethe*", Masson, Paris, 1983.

[4] L.D. Faddeev, E.K. Sklyanian and L.A. Takhtajan, The quantum inverse problem I, *Theor. Math. Phys.* **40** 194-220(1979).
P.P. Kulish, N.Yu Reshetikhin and E.K. Sklyanin, YBE and representations theory I, *Lett. Math. Phys.*, **5**(1981)393-403.

[5] P.P. Kulish and E.K. Sklyanin, Quantum spectral transformation method, recent developments, *Lecture Notes in Physics*, 151(Springer, 1982)pp61-119.
L.D. Faddeev, Quantum completely integrable models in field theory, *Math. Phys. Rev.*, ed. S.P. Novikov, Soviet scientific Reviews, Moskow, pp107-155, 1980.

[6] C.N. Yang, *Phys. Rev. Lett.* **19**(1967)1312-1314; *Phys. Rev.* **168**(1968)1920-1923.

[7] P.P. Kulish and E.K. Sklyanin, Solution to the YBE, *Zapicki Nauch. Semin. Otd. Matem. Inst. Steklova*, **95**(1980)129-160.

[8] R.J. Baxter, *Ann. Phys.*, **70**(1972)193-228; **76**(1973)1-24, **76**(1973)25-47;
"*Exactly solved models in statistical mechanics*", Academic, London(1982).

[9] H. Au-Yang, B.M. MaCoy, J.H.H. Perks, S. Tang and M.L. Yan, *Phys. Lett.* **A123**(1987)219;
R.J. Baxter, J.H.H. Perk and H. Au-Yang, *Phys. Lett.* **A128**(1988)668.

[10] V.G. Drinfeld, Hopf algebras and the Quantum Yang-Baxter equation, *Sov. Math. Dokl.* **32**, 254-258(1985).

[11] F.D.M. Haldane, *Phys. Rev. Lett.* **60**(1988)635;
B.S. Shastry, *Phys. Rev. Lett.* **60**(1988)639.

[12] D.B. Ugolov and V.E. Korepin, The Yangian sysmmetry of the Hubbard model, ITP-SB-93-66.

[13] F.D.M. Haldane, Z.N.C. Ha, J.C. Talstra, D. Bernard and V. Pasquier, *Phys. Rev. Lett.* **69**(1992)2021-2025.

[14] V. Chari and Pressley, Yangian and R-matrix, *L'Enseignement Mathematique*, **t36**(1990)267-302.

[15] F.D.M. Haldane, Physics of the ideal semion gas: spinons and quantum symmetries of the integrable Haldane-Shastry spin chain, in the Proceedings of the 16th Taniguchi Symposium on Condensed Matter Physics, 1993, edited by A. Okiji and N. Kawakami(Springer, 1994).

[16] F. Calogero, *J. Math. Phys.* **10**(1969)2191.
B. Sutherland, *J. Math. Phys.* **12**(1971)246; **12**(1971)251.
K. Hikami and M. Wadati, *J. Phys. Soc. Japan* **62**(1993)469; *Phys. Lett.* **A173**(1993)263.

[17] D. Bernard, M. Gaudin, F.D.M. Haldane and V. Pasquier, *J. Phys.* **A26** (1993)5219.

[18] A. Polychronakos, *Phys. Rev. Lett.* **69**(1992)703.
J.A. Minahan and A. Polychronakos, *Phys. Lett.* **B302**(1993)265.

[19] F. Calogero, O.Ragnisco and C. Marchiore, *Lett. Nuovo. Cimento* **13**(1975)383.

[20] B. Sutherland and R.A. Römer, *Phys. Rev. Lett.* **71**(1993)2789.

[21] M.L. Yan, Z. Chen, "Rational solution of Yang-Baxter equation and 1D many-body problem with generalized Delta-function", 1989, (unpublished).

[22] I.V. Komarov, The Goryachev-Chaplygin top in quantum mechanics, *Theor. Mat. Fiz.* **50**, 402(1982);
I.V. Komarov and V.V. Zalipaev, The Goryachev-Chaplygin Gyrostat in quantum mechanics, *J. Phys.* **A17**, 31(1984).

[23] E.K. Sklyanian, Quantum inverse scattering method , selected topics from "Quantum Group and Quantum Integrable Systems", World Scientific, Singapore(1991), pp.63-88.

[24] L.D. Faddeev, N.Yu Reshetikhin, L.A. Takhtajan, Quantization of Lie groups and Lie algebras, *Algebra and Analysis*, *1.1*, 118-206(1989)(in Russian).
N.Yu. Reshetikhin, M.A. Semenov-Tian-Shansky, Center extensions of quantum current groups, *Lett. Math. Phys.* **19**, 133-142(1990).

[25] V.G. Drinfeld, New realization of Yangian and quantum affine algebra, *Sov. Math. Dokl. 36*, 212-216(1988); Quantum Group, *Proc. of the ICM-86*, (Berkeley) Vol.1 New York. Academic press 1986, pp789-820.

[26] J. Ding and I.B. Frenkel, Isomorphism of two realizations of quantum affine algebra $U_q(\widehat{gl(n)})$, *Commun. Math. Phys.* **156**, 277-300(1993).

[27] M. Jimbo, "Introduction to the Yang-Baxter equation", Advanced series in mathematical physics, Vol.9, World Scientific(Singapore) 1989, pp.111-134.

[28] L.D. Faddeev, Quantum symmetry in Conformal Field Theory by Hamiltonian Methods, "Braid group, Knot theory and statistical mechanics II", eds. C.N. Yang and M.L. Ge, World Scientific, Singapore(1994), pp.108-129.

[29] J. Wess and B. Zumino, CERN preprint, CERN-TH-5697/90.

[30] M.L. Ge and K. Xue, Truncated $\widehat{SL}_q(2)$ algebra: Trigonometric Goryachev-Chaplygin Gyrostat, Nankai preprint(1994).